
Product Data Sheet

Pacific Blue[™] anti-mouse CD4

Catalog # / Size:	1102135 / 25 μg 1102140 / 100 μg
Clone:	GK1.5
Isotype:	Rat IgG2b, к
Immunogen:	Mouse CTL clone V4
Reactivity:	Mouse
Preparation:	The antibody was purified by affinity chromatography, and conjugated with Pacific Blue [™] under optimal conditions. The solution is free of unconjugated Pacific Blue [™] .
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration:	0.5



C57BL/6 mouse splenocytes were stained with CD4 (clone GK1.5) Pacfic BlueTM (filled histogram) or rat IgG2b, κ Pacific BlueTM isotype control (open histogram).

Applications:

Applications: Recommended Usage:	Flow Cytometry Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The suggested use of this reagent is \leq 1.0 microg per 10 ⁶ cells in 100 microL volume. It is highly recommended that the reagent be titrated for optimal performance for each application.
Application	* Pacific Blue [™] has a maximum emission of 455 nm when it is excited at 405 nm. Prior to using Pacific Blue [™] conjugate for flow cytometric analysis, please verify your flow cytometer's capability of exciting and detecting the fluorochrome.
Application Notes:	Additional reported applications (for the relevant formats) include: blocking of CD4 ⁺ T cell activation ^{1,4,11} , thymocyte costimulation3, <i>in vitro</i> and <i>in vivo</i> depletion ^{2,5-8} , blocking of egg-sperm cell adhesion ^{1,4} , immunohistochemical staining of acetone-fixed frozen sections ^{9,10} , and immunoprecipitation ^{1,2} . The GK1.5 antibody is able to block CD4 mediated cell adhesion and T cell activation. Binding of GK1.5 antibody to CD4 T cells can be blocked by RM4-5 antibody (Cat. No. 100506), but not RM4-4 antibody (Cat. No. 116002). The LEAF TM purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 100416). For <i>in vivo</i> studies or highly sensitive assays, we recommend Ultra-LEAF TM purified antibody (Cat. No. 100442) with a lower endotoxin limit than standard LEAF TM purified antibodies (Endotoxin <0.01 EU/microg).
Application References:	 Dialynas DP, <i>et al.</i> 1983. <i>J. Immunol.</i> 131:2445. (Block, IP) Dialynas DP, <i>et al.</i> 1983. <i>Immunol. Rev.</i> 74:29. (IP, Deplete) Wu L, <i>et al.</i> 1991. <i>J. Exp. Med.</i> 174:1617. (Costim) Godfrey DI, <i>et al.</i> 1994. <i>J. Immunol.</i> 152:4783. (Block) Gavett SH, <i>et al.</i> 1994. <i>Am. J. Respir. Cell. Mol. Biol.</i> 10:587. (Deplete) Schuyler M, <i>et al.</i> 1994. <i>Am. J. Respir. Crit. Care Med.</i> 149:1286. (Deplete) Gobrial RR, <i>et al.</i> 1989. <i>Clin. Immunol.</i> 142:954. (Deplete) Israelski DM, <i>et al.</i> 1986. <i>J. Immunol.</i> 184:1083. (IHC)

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com

10. Frei K, <i>et al.</i> 1997. <i>J. Exp. Med.</i> 185:2177. (IHC)	1
11. Felix NJ, <i>et al.</i> 2007. <i>Nat. Immunol.</i> 8:388. (Bloc	
12. Teo TH, <i>et al.</i> 2013. <i>J. Immunol.</i> 190:259. PubM	
13. Dekhtiarenko I, <i>et al.</i> 2013. <i>J. Immunol.</i> 190:33	99. <u>PubMed</u>
14. Morawski PA, <i>et al.</i> 2013. <i>J Biol Chem.</i> 288:244	94. <u>PubMed</u>
15. Wong LY, <i>et al.</i> 2013. <i>J Biol Chem.</i> 288:35170.	<u>PubMed</u>
16. Eitas TK, <i>et al.</i> 2014. <i>J Biol Chem.</i> 289:4173. <u>Pu</u>	<u>bMed</u>
17. Sawant DV, <i>et al.</i> 2014. <i>J. Immunol.</i> 192:2904.	
18. Madireddi S, <i>et al.</i> 2014. <i>J Exp Med.</i> 211:1433.	<u>PubMed</u>

Description:	CD4 is a 55 kD protein also known as L3T4 or T4. It is a member of the Ig superfamily, primarily expressed on most thymocytes, a subset of T cells, and weakly on macrophages and dendritic cells. It acts as a coreceptor with the TCR during T cell activation and thymic differentiation by binding MHC class II and associating with the protein tyrosin kinase, lck.
Antigen	1. Barclay A, <i>et al.</i> 1997. The Leukocyte Antigen FactsBook Academic Press.
References:	2. Bierer BE, <i>et al.</i> 1989. <i>Annu. Rev. Immunol.</i> 7:579.

3. Janeway CA. 1992. Annu. Rev. Immunol. 10:645.