
Product Data Sheet

KIRAVIA Blue 520™ anti-mouse CD4

Catalog # / Size: 1102390 / 100 µg
1102385 / 25 µg

Clone: GK1.5

Isotype: Rat IgG2b, κ

Immunogen: Mouse CTL clone V4

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with KIRAVIA Blue 520™ under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/mL

□ C57BL/6 mouse splenocytes were stained with CD3 APC and CD4 (clone GK1.5) KIRAVIA Blue 520™ (left) or rat IgG2b, κ KIRAVIA Blue 520™ isotype control (right).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 µg per million cells in 100 µL volume. It is recommended that the reagent be titrated for optimal performance for each application.

* KIRAVIA Blue 520™ has an excitation maximum of 495 nm, and a maximum emission of 520 nm.

Application Notes: Additional reported applications (for the relevant formats) include: blocking of CD4⁺ T cell activation^{1,4,11}, thymocyte costimulation³, *in vitro* and *in vivo* depletion^{2,5-8}, blocking of egg-sperm cell adhesion^{1,4}, immunohistochemical staining of acetone-fixed frozen sections^{9,10}, and immunoprecipitation^{1,2}. The GK1.5 antibody is able to block CD4 mediated cell adhesion and T cell activation. Binding of GK1.5 antibody to CD4 T cells can be blocked by RM4-5 antibody, but not RM4-4 antibody.

Application References:

1. Dialynas DP, *et al.* 1983. *J. Immunol.* 131:2445. (Block, IP)
2. Dialynas DP, *et al.* 1983. *Immunol. Rev.* 74:29. (IP, Deplete)
3. Wu L, *et al.* 1991. *J. Exp. Med.* 174:1617. (Costim)
4. Godfrey DI, *et al.* 1994. *J. Immunol.* 152:4783. (Block)
5. Gavett SH, *et al.* 1994. *Am. J. Respir. Cell. Mol. Biol.* 10:587. (Deplete)
6. Schuyler M, *et al.* 1994. *Am. J. Respir. Crit. Care Med.* 149:1286. (Deplete)
7. Ghobrial RR, *et al.* 1989. *Clin. Immunol. Immunopathol.* 52:486. (Deplete)
8. Israelski DM, *et al.* 1989. *J. Immunol.* 142:954. (Deplete)
9. Zheng B, *et al.* 1996. *J. Exp. Med.* 184:1083. (IHC)
10. Frei K, *et al.* 1997. *J. Exp. Med.* 185:2177. (IHC)
11. Felix NJ, *et al.* 2007. *Nat. Immunol.* 8:388. (Block)

Description: CD4 is a 55 kD protein also known as L3T4 or T4. It is a member of the Ig superfamily, primarily expressed on most thymocytes, a subset of T cells, and weakly on macrophages and dendritic cells. It acts as a coreceptor with the TCR during T cell activation and thymic differentiation by binding MHC class II and associating with the protein tyrosin kinase, lck.

Antigen
References:

1. Barclay A, *et al.* 1997. *The Leukocyte Antigen FactsBook* Academic Press.
2. Bierer BE, *et al.* 1989. *Annu. Rev. Immunol.* 7:579.
3. Janeway CA. 1992. *Annu. Rev. Immunol.* 10:645.