


Purified anti-mouse CD5

Catalog # / Size: 1103010 / 500 µg
Clone: 53-7.3
Isotype: Rat IgG2a, κ
Immunogen: Mouse thymus or spleen
Reactivity: Mouse
Preparation: The antibody was purified by affinity chromatography.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration: 0.5



C57BL/6 mouse splenocytes stained with purified 53-7.3, followed by anti-rat IgGs-FITC

Applications:

Applications: Other

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤1.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation¹, and immunohistochemistry² of acetone-fixed frozen tissue sections, zinc-fixed paraffin-embedded sections and formalin-fixed paraffin-embedded sections.

Application References:

- Ledbetter JA, *et al.* 1979. *Immunol. Rev.* 47:63. (IP)
- Ledbetter JA, *et al.* 1980. *J. Exp. Med.* 152:280. (FC, IHC)
- Bourdeau A, *et al.* 2007. *Blood* doi:10.1182/blood-2006-08-044370.

Description: CD5 is a 67 kD protein, also known as Lyt-1, Ly-1, T1, Tp67, or Ly-12. It is a member of the scavenger receptor cysteine-rich protein superfamily (SRCR) and primarily expressed on thymocytes, T cells, and B-1 cells. Although mature α/β T cells express high levels of CD5, very few γ/δ T cells express this antigen. The interaction of CD5 with CD72, gp35-37, TCR, or BCR is involved in T and B cell activation.

Antigen References:

- Barclay A, *et al.* 1997. *The Leukocyte Antigen FactsBook* Academic Press.
- Kipps TJ. 1988. *Adv. Immunol.* 47:117.
- Antin JH, *et al.* 1985. *J. Immunol.* 136:505.
- Tarakhovsky A, *et al.* 1