

**Brilliant Violet 510™ anti-mouse CD8a**

**Catalog # / Size:** 1103755 / 125 µl  
1103760 / 50 µg

**Clone:** 53-6.7

**Isotype:** Rat IgG2a, κ


**Immunogen:** Mouse thymus or spleen

**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 510™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 510™ and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

**Concentration:** microg sizes: 0.2 mg/ml  
microL sizes: lot-specific



C57BL/6 mouse splenocytes were stained with CD3 APC and CD8a (clone 53-6.7) Brilliant Violet 510™.

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining using the microg size, the suggested use of this reagent is ≤0.5 microg per million cells in 100 microL volume. For immunofluorescent staining using the microL size, the suggested use of this reagent is ≤5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 510™ excites at 405 nm and emits at 510 nm. The bandpass filter 510/50 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. **Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel.** Refer to your instrument manual or manufacturer for support. Brilliant Violet 510™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

**Application Notes:** Clone 53-6.7 antibody competes with clone 5H10-1 antibody for binding to thymocytes<sup>3</sup>. The 53-6.7 antibody has been reported to block antigen presentation via MHC class I and inhibit T cell responses to IL-2. This antibody has also been used for depletion of CD8a<sup>+</sup> cells. Additional reported applications (for the relevant formats) include: immunoprecipitation<sup>1,3</sup>, *in vivo* and *in vitro* cell depletion<sup>2,10,15</sup>, inhibition of CD8 T cell proliferation<sup>3</sup>, blocking of cytotoxicity<sup>3,4</sup>, and immunohistochemical staining<sup>5,6</sup> of acetone-fixed frozen sections and zinc-fixed paraffin-embedded sections. Clone 53-6.7 is not recommended for

immunohistochemistry of formalin-fixed paraffin sections. The LEAF™ purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 100716). For *in vivo* studies or highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 100746) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/microg).

**Application  
References:**

1. Ledbetter JA, *et al.* 1979. *Immunol. Rev.* 47:63. (IHC, IP)
2. Hathcock KS. 1991. *Current Protocols in Immunology.* 3.4.1. (Deplete)
3. Takahashi K, *et al.* 1992. *P. Natl. Acad. Sci. USA* 89:5557. (Block, IP)
4. Ledbetter JA, *et al.* 1981. *J. Exp. Med.* 153:1503. (Block)
5. Hata H, *et al.* 2004. *J. Clin. Invest.* 114:582. (IHC)
6. Fan WY, *et al.* 2001. *Exp. Biol. Med.* 226:1045. (IHC)
7. Shih FF, *et al.* 2006. *J. Immunol.* 176:3438. (FC)
8. Kamimura D, *et al.* 2006. *J. Immunol.* 177:306.
9. Bouwer HGA, *et al.* 2006. *P. Natl. Acad. Sci. USA* 103:5102. (FC, Deplete)
10. Kao C, *et al.* 2005. *Int. Immunol.* 17:1607. [PubMed](#)
11. Ko SY, *et al.* 2005. *J. Immunol.* 175:3309. (FC) [PubMed](#)
12. Rasmussen JW, *et al.* 2006. *Infect. Immun.* 74:6590. [PubMed](#)
13. Lee CH, *et al.* 2009. *Clin. Cancer Res.* [PubMed](#)
14. Geiben-Lynn R, *et al.* 2008. *Blood* 112:4585. (Deplete) [PubMed](#)
15. Kingeter LM, *et al.* 2008. *J. Immunol.* 181:6244. [PubMed](#)
16. Guo Y, *et al.* 2008. *Blood* 112:480. [PubMed](#)
17. Andrews DM, *et al.* 2008. *J. Virol.* 82:4931. [PubMed](#)
18. Britschqui MR, *et al.* 2008. *J. Immunol.* 181:7681. [PubMed](#)
19. Kenna TJ, *et al.* 2008. *Blood* 111:2091. [PubMed](#)
20. Jordan JM, *et al.* 2008. *Infect. Immun.* 76:3717. [PubMed](#)
21. Todd DJ, *et al.* 2009. *J. Exp. Med.* 206:2151. [PubMed](#)
22. Bankoti J, *et al.* 2010. *Toxicol. Sci.* 115:422. (FC) [PubMed](#)
23. Medyof H, *et al.* 2010. *Blood* 115:1175. [PubMed](#)
24. Riedl P, *et al.* 2009. *J. Immunol.* 183:370. [PubMed](#)
25. Apte SH, *et al.* 2010. *J. Immunol.* 185:998. [PubMed](#)
26. Bankoti J, *et al.* 2010. *Toxicol. Sci.* 115:422. (FC) [PubMed](#)
27. del Rio ML, *et al.* 2011. *Transpl. Int.* 24:501. (FC) [PubMed](#)
28. Wang T, *et al.* 2015. *PNAS.* 112:440. [PubMed](#)

---

**Description:**

CD8, also known as Lyt-2, Ly-2, or T8, consists of disulfide-linked  $\alpha$  and  $\beta$  chains that form the  $\alpha$ (CD8a)/ $\beta$ (CD8b) heterodimer and  $\alpha/\alpha$  homodimer. CD8a is a 34 kD protein that belongs to the immunoglobulin family. The CD8  $\alpha/\beta$  heterodimer is expressed on the surface of most thymocytes and a subset of mature TCR  $\alpha/\beta$  T cells. CD8 expression on mature T cells is non-overlapping with CD4. The CD8  $\alpha/\alpha$  homodimer is expressed on a subset of  $\gamma/\delta$  TCR-bearing T cells, NK cells, intestinal intraepithelial lymphocytes, and lymphoid dendritic cells. CD8 is an antigen co-receptor on T cells that interacts with MHC class I on antigen-presenting cells or epithelial cells. CD8 promotes T cell activation through its association with the TCR complex and protein tyrosine kinase lck.

**Antigen  
References:**

1. Barclay A, *et al.* 1997. *The Leukocyte Antigen FactsBook* Academic Press.
2. Zamoyska R. 1994. *Immunity* 1:243.
3. Ellmeier W, *et al.* 1999. *Annu. Rev. Immunol.* 17:523.