

PE anti-mouse CD23

Catalog # / Size: 1108035 / 50 µg
1108040 / 200 µg

Clone: B3B4

Isotype: Rat IgG2a, κ


Immunogen: Complex of IgE with Fcε receptor isolated from the mouse B hybridoma cell line O1.2B2

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2



CD57BL/6 mouse splenocytes were stained with CD23 (clone B3B4) PE (filled histogram), or rat IgG2a, κ PE (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 microg per 10⁶ cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for other applications.

Application Notes: The B3B4 antibody is useful for blocking IgE activity *in vivo*. Additional reported applications (for the relevant formats) include: immunoprecipitation¹, immunofluorescence microscopy, *in vitro* and *in vivo* blocking of ligand binding²⁻⁴, and immunohistochemical staining of acetone-fixed frozen sections⁵.

- Application References:**
1. Waldschmidt TJ, *et al.* 1988. *J. Immunol.* 140:2148. (IP)
 2. Rao M, *et al.* 1987. *J. Immunol.* 138:1845. (Block)
 3. Oshiba A, *et al.* 1997. *J. Immunol.* 159:4056. (Block)
 4. Dasic G, *et al.* 1999. *Eur. J. Immunol.* 29:2957. (Block)
 5. Maeda K, *et al.* 1992. *J. Immunol.* 148:2340. (IHC)
 6. Craig VJ, *et al.* 2011. *Cancer Res.* 71:3616. [PubMed](#)

Description: CD23 is a 45 kD protein also known as low affinity IgE Fc receptor, FcεRII, BLAST-2, Ly-42, or B6. It is a member of the Ig family, expressed on conventional B (but not B-1) cells and follicular dendritic cells. CD23 responds to high levels of IgE by downregulating IgE secretion.

- Antigen References:**
1. Barclay A, *et al.* 1997. *The Leukocyte Antigen FactsBook* Academic Press.
 2. Delespesse G, *et al.* 1992. *Immunol. Rev.* 125:77.
 3. Flores-Romo L, *et al.* 1993. *Science* 261:1038.