


Brilliant Violet 510™ anti-mouse CD23

Catalog # / Size: 1108115 / 50 µg
Clone: B3B4
Isotype: Rat IgG2a, κ
Immunogen: Complex of IgE with Fcε receptor isolated from the mouse B hybridoma cell line O1.2B2
Reactivity: Mouse
Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 510™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 510™ and unconjugated antibody.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).
Concentration: 0.2



C57BL/6 mouse splenocytes were stained with CD23 (clone B3B4) Brilliant Violet 510™ (filled histogram) or rat IgG2a, κ Brilliant Violet 510™ isotype control (open histogram).

Applications:

Applications: Flow Cytometry
Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.5 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 510™ excites at 405 nm and emits at 510 nm. The bandpass filter 510/50 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. **Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel.** Refer to your instrument manual or manufacturer for support. Brilliant Violet 510™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: The B3B4 antibody is useful for blocking IgE activity *in vivo*. Additional reported applications (for the relevant formats) include: immunoprecipitation¹, immunofluorescence microscopy, *in vitro* and *in vivo* blocking of ligand binding²⁻⁴, and immunohistochemical staining of acetone-fixed frozen sections⁵.

- Application References:**
1. Waldschmidt TJ, *et al.* 1988. *J. Immunol.* 140:2148. (IP)
 2. Rao M, *et al.* 1987. *J. Immunol.* 138:1845. (Block)
 3. Oshiba A, *et al.* 1997. *J. Immunol.* 159:4056. (Block)
 4. Dasic G, *et al.* 1999. *Eur. J. Immunol.* 29:2957. (Block)
 5. Maeda K, *et al.* 1992. *J. Immunol.* 148:2340. (IHC)

Description: CD23 is a 45 kD protein also known as low affinity IgE Fc receptor, FcεRII, BLAST-2, Ly-42, or B6. It is a member of the Ig family, expressed on conventional B (but not B-1) cells and follicular dendritic cells. CD23 responds to high levels of IgE by downregulating IgE secretion.

Antigen
References:

1. Barclay A, *et al.* 1997. *The Leukocyte Antigen FactsBook* Academic Press.
2. Delespesse G, *et al.* 1992. *Immunol. Rev.* 125:77.
3. Flores-Romo L, *et al.* 1993. *Science* 261:1038.