SONY

Product Data Sheet

PE/Cyanine7 anti-mouse CD24

Catalog # / Size: 1109105 / 25 µg
1109110 / 100 µg

Clone: M1/69

Isotype: Rat IgG2b, κ

Immunogen: C57BL/10 mouse splenic T cells and concanavalin A-activated splenocytes

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE/Cy7 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2



C57BL/6 mouse splenocytes stained with M1/69 PE/Cy7

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 µg per 10^6 cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: Western blotting1, *in vitro* induction of thymocyte maturation2, complement-mediated cytotoxicity3, and immunohistochemistry of acetone-fixed frozen sections4, formalin-fixed paraffin-embedded sections5 and zinc-fixed paraffin-embedded sections[10]. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 101810).

Application References:

1. Springer T, *et al.* 1978. *Eur. J. Immunol.* 8:539. (WB)
2. Crowley M, *et al.* 1989. *Cell. Immunol.* 118:108. (FA)
3. Veillette A, *et al.* 1989. *J. Exp. Med.* 170:1671. (FA)
4. Pandelakis A Flavell RA 1999 *JEM* 189:855 (FC, IHC)
5. Liu JQ, *et al.* 2007 *J. Immunol.* 178:6227. (FC, IF)
6. Chappaz S, *et al.* 2007. *Blood* doi:10.1182/blood-2007-02-074245. (FC) PubMed
7. Rucci F, *et al.* 2010. *Proc Natl Acad Sci USA.* 107:3024. (FC) PubMed
8. Teague TK, *et al.* 2010. *Int Immunol.* 22:387. (FC) PubMed
9. Gracz AD, *et al.* 2010. *Am J. Physiol Gastrointest Liver Physiol.* 298:590. (FC) PubMed
10. Chen CY, *et al.* 2008. *Endocrinology.* 10:1210. (FC, IHC) PubMed
11. Qui Q, *et al.* 2010. *J. Immunol.* 184:1681. (FC) PubMed

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.

Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com

Description: CD24 is a 35-45 kD protein also known as Heat Stable Antigen (HSA), Ly-52, or Nectadrin. It is a GPI-linked sialoglycoprotein expressed on lymphocytes, granulocytes, epithelial cells, thymocytes, monocytes, erythrocytes, and dendritic cells. CD24 expression varies during T and B cell differentiation and is a useful marker for delineating various lymphocyte developmental stages. CD24 serves as an adhesion or costimulatory molecule involved in T and B lymphocyte activation and differentiation by homophilic binding or binding to CD62P.

Antigen References:

1. Barclay A, *et al.* 1997. The Leukocyte Antigen FactsBook Academic Press.
2. Aigner S, *et al.* 1997. *Blood* 89:3385.
3. Hough MR, *et al.* 1996. *J. Immunol.* 156:479.
4. Liu Y, *et al.* 1992. *J. Exp. Med.* 175:437.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.