
## **Product Data Sheet**

## PerCP/Cy5.5 anti-mouse CD25

Catalog # / Size:	1109555 / 25 μg 1109560 / 100 μg
Clone:	3C7
Isotype:	Rat IgG2b, к
Immunogen:	IL-2-dependent BALB/c mouse helper T- cell clone HT-2
<b>Reactivity:</b>	Mouse
Preparation:	The antibody was purified by affinity chromatography and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
<b>Concentration:</b>	0.2



Con A-stimulated C57BL/7 splenocytes (Day 3) stained with 3C7 PerCP/Cy5.5

## **Applications:**

Applications:	Flow Cytometry
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤1.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application. * PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.
Application Notes:	Additional reported applications (for the relevant formats) include: <i>in vitro</i> blocking of IL-2 binding to low- and high-affinity receptors <sup>1,2</sup> , and immunohistochemical staining of acetone-fixed frozen sections. 3C7 antibody recognizes different epitope of PC61 antibody (Cat. No. 102002). The LEAF <sup>™</sup> Purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 101906).
Application References:	1. Ortega RG, <i>et al.</i> 1984. <i>J. Immunol.</i> 133:1970. (Block) 2. Moreau JL, <i>et al.</i> 1987. <i>Eur. J. Immunol.</i> 17:929. (Block) 3. Lei F, <i>et al.</i> 2013. <i>J Vis Exp.</i> 60:3986. <u>PubMed.</u>
Description:	CD25 is a 55 kD glycoprotein, also known as the low affinity IL-2R $\alpha$ , Ly-43, p55, or Tac. It is expressed on activated T and B cells, thymocyte subset, pre-B cells, and T regulatory cells. In association with CD122 (IL-2R $\beta$ ) and CD132(common $\gamma$ chain), CD25 forms the high affinity signaling IL-2 receptor.
Antigen References:	1. Taniguchi T, <i>et al.</i> 1993. <i>Cell</i> 73:5. 2. Waldmann TA. 1991. <i>J. Biol. Chem.</i> 266:2681. 3. Read S, <i>et al.</i> 2000. <i>J. Exp. Med.</i> 192:295. 4. Lowenthal JW, <i>et al.</i> 1985. <i>J. Immunol.</i>

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com