SONY

Product Data Sheet

PerCP/Cyanine5.5 anti-mouse CD31

Catalog # / Size: 1112610 / 100 µg

Clone: MEC13.3

Isotype: Rat IgG2a, κ

Immunogen: Polyoma middle T transformed EC line tEnd.1

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with PerCP/Cyanine5.5 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Workshop Number: 750 under optimal conditions.

Concentration: 0.2



C57BL/6 mouse splenocytes were stained with CD31 (clone MEC13.3) PerCp/Cyanine5.5 (filled histogram) or Rat IgG2a, κ PerCp/Cyanine5.5 isotype control (open histogram)

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

* PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

Application Notes: Anti-mouse CD31 clones 390 and MEC13.3 bind to their respective non-overlapping epitopes in IgD2 of CD31.[8] Additional reported applications (in the relevant formats) include: immunoprecipitation1, *in vitro* and *in vivo* blocking of CD31-mediated cell-cell interactions[1-4], and immunohistochemical staining[1,5,6] of acetone-fixed frozen sections and zinc-fixed paraffin-embedded sections.
Special Note: The antibody works well on acetone-fixed frozen sections as well as Zinc-fixed paraffin-embedded sections. It sometime works on formalin-fixed and paraformaldehyde-fixed paraffin-embedded tissue sections but inconsistent results have been reported. This antibody is not recommended for formalin-fixed paraffin-embedded sections or for Western blot analysis. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 102512).

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.

Application References:

1. Vecchi A, *et al.* 1994. *Eur. J. Cell Biol.* 63:247. (IP, IHC, Block)
2. Christofidou-Solomidou M, *et al.* 1997. *J. Immunol.* 158:4872. (Block)
3. DeLisser HM, *et al.* 1997. *Am. J. Pathol.* 151:671. (Block)
4. Rosenblum WI, *et al.* 1994. *Am. J. Pathol.* 145:33. (Block)
5. Baldwin HS, *et al.* 1994. *Development* 120:2539. (IHC)
6. Voswinckel R, *et al.* 2003. *Circ. Res.* 93:372. (IHC)
7. Leung VW, *et al.* 2009. *Am J. Pathol.* 175:1757. PubMed
8. Chacko AM, *et al.* 2012. *PLoS One* 7:e34958.
9. Giacomini C, *et al.* 2014. *Exp Eye Res.* 18:1. PubMed
10. Morita R, *et al.* 2015. *PNAS.* 112:160. PubMed
11. Ito A, *et al.* 2015. *Brain Res.* 1594:310. PubMed

Description:

CD31 is a 130-140 kD glycoprotein, also known as platelet endothelial cell adhesion molecule (PECAM-1), EndoCAM, and gpIIa. It is a member of the Ig superfamily, expressed on endothelial cells, platelets, granulocytes, monocytes/macrophages, dendritic cells, and T and B cell subsets, and is critical for cell-to-cell interactions. The primary ligands for CD31 have been reported to be CD38 and the vitronectin receptor (α_v β_3 integrin, CD51/CD61). Other reported functions of CD31 are neutrophil emigration to sites of inflammation, and angiogenesis.

Antigen References:

1. Barclay AN, *et al.* 1997. The Leukocyte Antigen FactsBook Academic Press.
2. DeLisser HM, *et al.* 1994. *Immunol. Today* 15:490.
3. Newman PJ, *et al.* 1990. *Science* 247:1219.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.