

APC/Fire™ 750 anti-mouse CD31

Catalog # / Size: 1112635 / 25 µg
1112640 / 100 µg

Clone: MEC13.3

Isotype: Rat IgG2a, κ

Immunogen: Polyoma middle T transformed EC line tEnd.1


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 750 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Workshop Number: 750 under optimal conditions.

Concentration: 0.2 mg/ml




C57BL/6 were stained with CD31 (clone MEC13.3) APC/Fire™ 750 (filled histogram) or rat IgG2a, κ APC/Fire™ 750 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

* APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 787 nm.



Application Notes: Anti-mouse CD31 clones 390 and MEC13.3 bind to their respective non-overlapping epitopes in IgD2 of CD31.⁸ Additional reported applications (in the relevant formats) include: immunoprecipitation¹, *in vitro* and *in vivo* blocking of CD31-mediated cell-cell interactions¹⁻⁴, and immunohistochemical staining^{1,5,6} of acetone-fixed frozen sections and zinc-fixed paraffin-embedded sections.

Special Note: The antibody works well on acetone-fixed frozen sections as well as Zinc-fixed paraffin-embedded sections. It sometime works on formalin-fixed and paraformaldehyde-fixed paraffin-embedded tissue sections but inconsistent results have been reported. This antibody is not recommended for formalin-fixed paraffin-embedded sections or for Western blot analysis. The Ultra-LEAF™ Purified antibody (Endotoxin < 0.01 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 102529 and 102530).

Application References:

1. Vecchi A, *et al.* 1994. *Eur. J. Cell Biol.* 63:247. (IP, IHC, Block)
2. Christofidou-Solomidou M, *et al.* 1997. *J. Immunol.* 158:4872. (Block)
3. DeLisser HM, *et al.* 1997. *Am. J. Pathol.* 151:671. (Block)
4. Rosenblum WI, *et al.* 1994. *Am. J. Pathol.* 145:33. (Block)
5. Baldwin HS, *et al.* 1994. *Development* 120:2539. (IHC)
6. Voswinckel R, *et al.* 2003. *Circ. Res.* 93:372. (IHC)
7. Leung VW, *et al.* 2009. *Am J. Pathol.* 175:1757. [PubMed](#)
8. Chacko AM, *et al.* 2012. *PLoS One* 7:e34958.
9. Giacomini C, *et al.* 2014. *Exp Eye Res.* 18:1. [PubMed](#)
10. Morita R, *et al.* 2015. *PNAS.* 112:160. [PubMed](#)
11. Ito A, *et al.* 2015. *Brain Res.* 1594:310. [PubMed](#)

Description: CD31 is a 130-140 kD glycoprotein, also known as platelet endothelial cell adhesion molecule (PECAM-1), EndoCAM, and gpIIa. It is a member of the Ig superfamily, expressed on endothelial cells, platelets, granulocytes, monocytes/macrophages, dendritic cells, and T and B cell subsets, and is critical for cell-to-cell interactions. The primary ligands for CD31 have been reported to be CD38 and the vitronectin receptor ($\alpha_v \beta_3$ integrin, CD51/CD61). Other reported functions of CD31 are neutrophil emigration to sites of inflammation, and angiogenesis.

Antigen References:

1. Barclay AN, *et al.* 1997. The Leukocyte Antigen FactsBook Academic Press.
2. DeLisser HM, *et al.* 1994. *Immunol. Today* 15:490.
3. Newman PJ, *et al.* 1990. *Science* 247:1219.