

APC anti-mouse CD38

Catalog # / Size: 1113560 / 100 µg
1113555 / 25 µg

Clone: 90

Isotype: Rat IgG2a, κ


Immunogen: Mouse bone marrow pre-B cells

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2



C57BL/6 mouse splenocytes stained with 90 APC

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemistry^{1,2} of acetone-fixed frozen sections, and induction of B cell proliferation¹.

Application References: 1. Oliver AM, *et al.* 1997. *J. Immunol.* 158:1108.
2. Howard M, *et al.* 1993. *Science* 262:1056.

Description: CD38 is a 42 kD glycoprotein, also known as T10. It is an ADP-ribosyl hydrolase, expressed on B cells, NK cells, a subset of T cells, brain, muscle, and kidney. In mouse, CD38 expression is downregulated on germinal center B cells and plasma cells, whereas this is not the case for humans. By functioning as both a cyclase and a hydrolase, CD38 mediates lymphocyte activation, as well as adhesion and metabolism of cADPR and NAADP. CD31 is the ligand of CD38.

Antigen References: 1. Barclay AN, *et al.* 1997. The Leukocyte Antigen FactsBook Academic Press.
2. Shubinsky G, *et al.* 1997. *Immunity* 7:315.
3. Cesano A, *et al.* 1998. *J. Immunol.* 160:1106.
4. Cockayne DA,