
Product Data Sheet

Brilliant Violet 605[™] anti-mouse CD45

Catalog # / Size:	1115700 / 500 μl 1115695 / 125 μl	
	1115775 / 50 μg	
Clone:	30-F11	umbe
Isotype:	Rat IgG2b, к	Relative Cell Numbe
Immunogen:	Mouse thymus or spleen	ative (
Reactivity:	Mouse	B
Preparation:	The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 605 [™] under optimal conditions. The solution is free of unconjugated Brilliant Violet 605 [™] and unconjugated antibody.	C5 sta
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).	Br his Vic
Concentration:	microg sizes: 0.2 mg/ml microL sizes: lot-specific	his



C57BL/6 mouse splenocytes were stained with CD45 (clone 30-F11) Brilliant Violet 605[™] (filled histogram) or rat IgG2b, κ Brilliant Violet 605[™] isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining with microg, the suggested use of this reagent is ≤0.5 microg per million cells in 100 microL volume. For flow cytometric staining with microL, the suggested use of this reagent is ≤5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 605[™] excites at 405 nm and emits at 603 nm. The bandpass filter 610/20 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. **Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel.** Refer to your instrument manual or manufacturer for support. Brilliant Violet 605[™] is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

ApplicationClone 30-F11 reacts with all isoforms and both CD45.1 and CD45.2 alloantigens of
CD45.Notes:CD45.

Additional reported applications (for relevant formats) include: immunoprecipitation3, complement-dependent cytotoxicity^{1,5}, immunohistochemistry (acetone-fixed frozen sections, zinc-fixed paraffinembedded sections and formalin-fixed paraffin-embedded sections)^{4,6} and

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com Western blotting⁷. The LEAF[™] purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 103120).
Application
1. Podd BS, *et al.* 2006. *J. Immunol.* 176:6532. (FC, CMCD) <u>PubMed</u>
2. Haynes NM, *et al.* 2007. *J. Immunol.* 179:5099. (FC)
3. Ledbetter JA, *et al.* 1979. *Immunol. Rev.* 47:63. (IP)
4. Simon DI, *et al.* 2000. *J. Clin. Invest.* 105:293. (IHC)
5. Seaman WE. 1983. *J. Immunol.* 130:1713. (CMCD)
6. Cornet A, *et al.* 2001. *P. Natl. Acad. Sci. USA* 98:13306. (IHC)
7. Tsuboi S and Fukuda M. 1998. *J. Biol. Chem.* 273:30680. (WB) <u>PubMed</u>
8. Liu F, *et al.* 2012. *Blood.* 119:3295. <u>PubMed</u>
9. Pelletier AN, *et al.* 2012. *J. Immunol.* 188:5561. <u>PubMed</u>

Description:	CD45 is a 180-240 kD glycoprotein also known as the leukocyte common antigen (LCA), T200, or Ly-5. It is a member of the protein tyrosine phosphatase (PTP) family, expressed on all hematopoietic cells except mature erythrocytes and platelets. There are different isoforms of CD45 that arise from variable splicing of exons 4, 5, and 6, which encode A, B, and C determinants, respectively. CD45 plays a key role in TCR and BCR signal transduction. These isoforms are very specific to the activation and maturation state of the cell as well as cell type. The primary ligands for CD45 are galectin-1, CD2, CD3, CD4, TCR, CD22, and Thy-1.

Antigen	1. Barclay A, <i>et al.</i> 1997. The Leukocyte Antigen FactsBook Academic Press.
References:	2. Trowbridge IS, et al. 1993. Annu. Rev. Immunol. 12:85.
	3. Kishihara K, <i>et al.</i> 1993. <i>Cell</i> 74:143.
	4. Pulido R, <