

**Alexa Fluor® 647 anti-mouse/human CD45R/B220**

**Catalog # / Size:** 1116130 / 100 µg  
1116145 / 25 µg

**Clone:** RA3-6B2

**Isotype:** Rat IgG2a, κ


**Immunogen:** Abelson murine leukemia virus-induced pre-B tumor cells

**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with Alexa Fluor® 647 under optimal conditions.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.5




C57BL/6 mouse splenocytes stained with RA3-6B2 Alexa Fluor® 647

**Applications:**

**Applications:** Immunofluorescence

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. For immunohistochemistry, a concentration range of 2.5-5 microg/ml is suggested. For immunofluorescence microscopy, a concentration range of 1.25-10 microg/ml is recommended. It is recommended that the reagent be titrated for optimal performance for each application.



C57BL/6 mouse frozen lymph node sections were fixed with 4% paraformaldehyde (PFA) for 10 minutes at room temperature and blocked with 5% FBS plus 5% rat serum for 1 hour at room temperature. Then the section was stained with 5 microg/ml of CD4 (clone G)

\* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633nm / 635nm.

**Application Notes:** Clone RA3-6B2 has been described to react with an epitope on the extracellular domain of the transmembrane CD45 glycoprotein which is dependent upon the expression of exon A and specific carbohydrate residues. Additional reported applications (for the relevant formats) include: immunoprecipitation<sup>1</sup>, *in vitro* and *in vivo* modulation of B cell responses<sup>2-4</sup>, and immunohistochemistry of acetone-fixed frozen sections and formalin-fixed paraffin-embedded sections<sup>5,6</sup>. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered)

is recommended for functional assays  
(Cat. No. 103216).

- Application** 1. Coffman RL. 1982. *Immunol. Rev.* 69:5. (IP)
- References:** 2. George A, *et al.* 1994. *J. Immunol.* 152:1014. (Activ)  
3. Asensi V, *et al.* 1989. *Immunology* 68:204. (Activ)  
4. Domiati-Saad R, *et al.* 1993. *J. Immunol.* 151:5936. (Activ)  
5. Hata H, *et al.* 2004. *J. Clin. Invest.* 114:582. (IHC)  
6. Monteith CE, *et al.* 1996. *Can. J. Vet. Res.* 60:193. (IHC)  
7. Shih FF, *et al.* 2006. *J. Immunol.* 176:3438. (FC)  
8. Chang C L-T, *et al.* 2007. *J. Immunol.* 178:6984.  
9. Fazilleau N, *et al.* 2007. *Nature Immunol.* 8:753.  
10. Lang GL, *et al.* 2008. *Blood* 111:2158. [PubMed](#)  
11. Charles N, *et al.* 2010. *Nat. Med.* 16:701. (FC) [PubMed](#)  
12. del Rio ML, *et al.* 2011. *Transpl. Int.* 24:501. (FC) [PubMed](#)  
13. Murakami R, *et al.* 2013. *PLoS One.* 8:73270. [PubMed](#)
- 

**Description:** CD45R, also known as B220, is an isoform of CD45. It is a member of the protein tyrosine phosphatase (PTP) family with a molecular weight of approximately 180-240 kD. CD45R is expressed on B cells (at all developmental stages from pro-B cells through mature B cells), activated B cells, and subsets of T and NK cells. CD45R (B220) is also expressed on a subset of abnormal T cells involved in the pathogenesis of systemic autoimmunity in MRL-*Fas<sup>lpr</sup>* and MRL-*Fas<sup>gld</sup>* mice. It plays a critical role in TCR and BCR signaling. The primary ligands for CD45 are galectin-1, CD2, CD3, and CD4. CD45R is commonly used as a pan-B cell marker; however, CD19 may be more appropriate for B cell specificity.

- Antigen** 1. Barclay A, *et al.* 1997. *The Leukocyte Antigen FactsBook* Academic Press.
- References:** 2. Trowbridge IS, *et al.* 1993. *Annu. Rev. Immunol.* 12:85.  
3. Kishihara K, *et al.* 1993. *Cell* 74:143.  
4. Pulido R, <