


Purified anti-mouse CD45RB

Catalog # / Size: 1116510 / 500 µg
Clone: C363-16A
Isotype: Rat IgG2a, κ
Immunogen: Cloned mouse Th2 cell lines
Reactivity: Mouse
Preparation: The antibody was purified by affinity chromatography.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration: 0.5



C57BL/6 mouse splenocytes stained with purified C363-16A, followed by anti-rat IgG FITC

Applications:

Applications: Other

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.06 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation¹, immunohistochemistry of acetone-fixed frozen sections and formalin-fixed paraffin-embedded sections.

Application References: 1. Bottomly K, *et al.* 1989. *Eur. J. Immunol.* 19:617.
2. Norian LA and Allen PM. 2004. *J. Immunol.* 173:835.

Description: CD45RB is an isoform of CD45 with exon 5 splicing (encodes B determinant). It is a 220 kD glycoprotein expressed on peripheral B cells, naïve T cells, thymocytes, weakly on macrophages, and dendritic cells. It plays a critical role in TCR and BCR signaling. As T cells become activated and progress from naïve to memory cells, CD45RB expression is downregulated. Additionally, functionally distinct CD4⁺ T cell subsets, which secrete differing cytokine profiles, can be separated by CD45RB intensity. The primary ligands for CD45 are galectin-1, CD2, CD3, CD4 and Thy-1.

Antigen References: 1. Barclay A, *et al.* 1997. The Leukocyte Antigen FactsBook Academic Press.
2. Trowbridge IS, *et al.* 1993. *Annu. Rev. Immunol.* 12:85.
3. Kishihara K, *et al.* 1993. *Cell* 74:143.
4. Pulido R, <