

PerCP/Cy5.5 anti-mouse CD45RB

Catalog # / Size: 1116570 / 100 µg
1116565 / 25 µg

Clone: C363-16A

Isotype: Rat IgG2a, κ


Immunogen: Cloned mouse Th2 cell lines

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2



C57BL/6 mouse splenocytes stained with C363-16A PerCP/Cy5.5

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 microg per 10⁶ cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

* PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

- Application References:**
1. Bottomly K, *et al.* 1989. *Eur. J. Immunol.* 19:617.
 2. Norian LA and Allen PM. 2004. *J. Immunol.* 173:835.

Description: CD45RB is an isoform of CD45 with exon 5 splicing (encodes B determinant). It is a 220 kD glycoprotein expressed on peripheral B cells, naïve T cells, thymocytes, weakly on macrophages, and dendritic cells. It plays a critical role in TCR and BCR signaling. As T cells become activated and progress from naïve to memory cells, CD45RB expression is downregulated. Additionally, functionally distinct CD4⁺ T cell subsets, which secrete differing cytokine profiles, can be separated by CD45RB intensity. The primary ligands for CD45 are galectin-1, CD2, CD3, CD4 and Thy-1.

- Antigen References:**
1. Barclay A, *et al.* 1997. *The Leukocyte Antigen FactsBook* Academic Press.
 2. Trowbridge IS, *et al.* 1993. *Annu. Rev. Immunol.* 12:85.
 3. Kishihara K, *et al.* 1993. *Cell* 74:143.
 4. Pulido R, *et al.* <