

Pacific Blue™ anti-mouse CD45RB

Catalog # / Size: 1116575 / 25 µg
1116580 / 100 µg

Clone: C363-16A

Isotype: Rat IgG2a, κ


Immunogen: Cloned mouse Th2 cell lines

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with Pacific Blue™ under optimal conditions. The solution is free of unconjugated Pacific Blue™.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5



C57BL/6 splenocytes stained with C363-16A Pacific Blue™

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The suggested use of this reagent is ≤ 1.0 microg per 10^6 cells in 100 microL volume or 100 microL of whole blood. It is highly recommended that the reagent be titrated for optimal performance for each application.

* Pacific Blue™ has a maximum emission of 455 nm when it is excited at 405 nm. Prior to using Pacific Blue™ conjugate for flow cytometric analysis, please verify your flow cytometer's capability of exciting and detecting the fluorochrome.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation¹, immunohistochemistry of acetone-fixed frozen sections and formalin-fixed paraffin-embedded sections.

Application References: 1. Bottomly K, *et al.* 1989. *Eur. J. Immunol.* 19:617.
2. Norian LA and Allen PM. 2004. *J. Immunol.* 173:835.

Description: CD45RB is an isoform of CD45 with exon 5 splicing (encodes B determinant). It is a 220 kD glycoprotein expressed on peripheral B cells, naïve T cells, thymocytes, weakly on macrophages, and dendritic cells. It plays a critical role in TCR and BCR signaling. As T cells become activated and progress from naïve to memory cells, CD45RB expression is downregulated. Additionally, functionally distinct CD4⁺ T cell subsets, which secrete differing cytokine profiles, can be separated by CD45RB intensity. The primary ligands for CD45 are galectin-1, CD2, CD3, CD4 and Thy-1.

Antigen References: 1. Barclay A, *et al.* 1997. *The Leukocyte Antigen FactsBook* Academic Press.
2. Trowbridge IS, *et al.* 1993. *Annu. Rev. Immunol.* 12:85.
3. Kishihara K, *et al.* 1993. *Cell* 74:143.
4. Pulido R,