


Purified anti-mouse CD48

Catalog # / Size: 1117010 / 500 µg
Clone: HM48-1
Isotype: Hamster IgG
Immunogen: Mouse T lymphoma MBL-2
Reactivity: Mouse
Preparation: The antibody was purified by affinity chromatography.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration: 0.5




C57BL/6 mouse splenocytes stained with purified HM48-1, followed by anti-Armenian hamster IgG FITC.

Applications:

Applications: Other

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 microg per 10⁶ cells in 100 microL volume. For immunohistochemical staining of frozen sections, a concentration range of 1.0 - 5.0 microg/ml is suggested. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: The HM48-1 antibody is useful for blocking *in vitro* and *in vivo* CD48 mediated interactions. Additional reported applications (for the relevant formats) include: immunoprecipitation¹, costimulation of T cell proliferation^{1,2}, blocking of CD48-CD2 interaction¹, and inhibition of CTL activity and graft rejection^{1,2}. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 103408). For *in vivo* studies or highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 103430) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/microg).



C57BL/6 mouse frozen spleen section was fixed with 4% paraformaldehyde (PFA) for ten minutes at room temperature and blocked with 5% rat serum + FBS for one hour at room temperature. Then the section was stained with 5 µg/ml of purified CD48 (clone HM4)

Application 1. Kato K, *et al.* 1992. *J. Exp. Med.* 176:1241. (IP, Costim, Block)

References: 2. Qin L, *et al.* 1994. *J. Exp. Med.* 179:341. (Costim, Block)

Description: CD48 is a 45 kD GPI-anchored glycoprotein also known as BCM1, Blast-1 (human), and OX-45 (rat). It is a member of the Ig superfamily, expressed on T and B cells and monocytes/macrophages. It plays a role in adhesion and T cell recognition. The primary ligands for CD48 are CD2 and CD244.

Antigen
References: 1. Barclay AN, *et al.* 1997. *The Leukocyte Antigen FactsBook* Academic Press.
2. Flament C, *et al.* 1996. *Hum. Immunol.* 46:82.
3. Van der Merwe PA, *et al.* 1995. *Curr. Biol.* 5:74.
4. Latchman Y