PE/Cy5 anti-mouse CD48

Catalog # / Size: 1117100 / 100 μg

Clone: HM48-1
Isotype: Hamster IgG

Immunogen: Mouse T lymphoma MBL-2

Reactivity: Mouse

Preparation: The antibody was purified by affinity


chromatography, and conjugated with PE/Cy5 under optimal conditions. The solution is free of unconjugated PE/Cy5

and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

Concentration: 0.2



C57BL/6 splenocytes stained with HM48-1 PE/CY5

Applications:

Applications: Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of

this reagent is \leq 0.25 microg per 10^6 cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each

application.

Application

Notes:

The HM48-1 antibody is useful for blocking *in vitro* and *in vivo* CD48 mediated interactions. Additional reported applications (for the relevant formats) include: immunoprecipitation1, costimulation of T cell proliferation 1,2 , blocking of CD48-CD2 interaction1, and inhibition of CTL activity and graft rejection 1,2 . The LEAF $^{\text{TM}}$ purified antibody (Endotoxin <0.1 EU/ μ g, Azide-Free, 0.2 μ m filtered) is recommended for functional assays (Cat. No. 103408). For *in vivo* studies or highly sensitive assays, we recommend Ultra-LEAF $^{\text{TM}}$ purified antibody (Cat. No. 103430) with a lower endotoxin limit than standard LEAF $^{\text{TM}}$ purified antibodies (Endotoxin <0.01 EU/microg).

Application References:

1. Kato K, et al. 1992. J. Exp. Med. 176:1241. (IP, Costim, Block) 2. Qin L, et al. 1994. J. Exp. Med. 179:341. (Costim, Block)

Description:

CD48 is a 45 kD GPI-anchored glycoprotein also known as BCM1, Blast-1 (human), and OX-45 (rat). It is a member of the lg superfamily, expressed on T and B cells and monocytes/macrophages. It plays a role in adhesion and T cell recognition. The primary ligands for CD48 are CD2 and CD244.

Antigen References:

1. Barclay AN, et al. 1997. The Leukocyte Antigen FactsBook Academic Press.

2. Flament C, et al. 1996. Hum. Immunol. 46:82.

3. Van der Merwe PA, et al. 1995. Curr. Biol. 5:74.

4. Latchman Y