

## **Product Data Sheet**

## PE/Cy7 anti-mouse CD48

Catalog # / Size:	1117115 / 25 μg 1117120 / 100 μg
Clone:	HM48-1
Isotype:	Hamster IgG
Immunogen:	Mouse T lymphoma MBL-2
<b>Reactivity:</b>	Mouse
Preparation:	The antibody was purified by affinity chromatography, and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
<b>Concentration:</b>	0.2





## **Applications:**

Applications:	Flow Cytometry
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 0.06$ microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.
Application Notes:	The HM48-1 antibody is useful for blocking <i>in vitro</i> and <i>in vivo</i> CD48 mediated interactions. Additional reported applications (for the relevant formats) include: immunoprecipitation1, costimulation of T cell proliferation <sup>1,2</sup> , blocking of CD48-CD2 interaction1, and inhibition of CTL activity and graft rejection <sup>1,2</sup> . The LEAF <sup>™</sup> purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 103408). For <i>in vivo</i> studies or highly sensitive assays, we recommend Ultra-LEAF <sup>™</sup> purified antibody (Cat. No. 103430) with a lower endotoxin limit than standard LEAF <sup>™</sup> purified antibodies (Endotoxin <0.01 EU/microg).
Application References:	1. Kato K, <i>et al.</i> 1992. <i>J. Exp. Med.</i> 176:1241. (IP, Costim, Block) 2. Qin L, <i>et al.</i> 1994. <i>J. Exp. Med.</i> 179:341. (Costim, Block)
Description:	CD48 is a 45 kD GPI-anchored glycoprotein also known as BCM1, Blast-1 (human), and OX-45 (rat). It is a member of the Ig superfamily, expressed on T and B cells and monocytes/macrophages. It plays a role in adhesion and T cell recognition. The primary ligands for CD48 are CD2 and CD244.
Antigen References:	<ol> <li>Barclay AN, <i>et al.</i> 1997. The Leukocyte Antigen FactsBook Academic Press.</li> <li>Flament C, <i>et al.</i> 1996. <i>Hum. Immunol.</i> 46:82.</li> <li>Van der Merwe PA, <i>et al.</i> 1995. <i>Curr. Biol.</i> 5:74.</li> <li>Latchman Y</li> </ol>

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com