


PE/Dazzle™ 594 anti-mouse CD48

Catalog # / Size:	1117190 / 100 µg 1117185 / 25 µg
Clone:	HM48-1
Isotype:	Hamster IgG
Immunogen:	Mouse T lymphoma MBL-2
Reactivity:	Mouse
Preparation:	The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions. The solution is free of unconjugated PE/Dazzle™ 594 and unconjugated antibody.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration:	0.2



C57BL/6 mouse splenocytes were stained with CD48 (clone HM48-1) PE/Dazzle™ 594 (filled histogram) or Armenian hamster IgG PE/Dazzle™ 594 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.

Application Notes: The HM48-1 antibody is useful for blocking *in vitro* and *in vivo* CD48 mediated interactions. Additional reported applications (for the relevant formats) include: immunoprecipitation¹, costimulation of T cell proliferation^{1,2}, blocking of CD48-CD2 interaction¹, and inhibition of CTL activity and graft rejection^{1,2}. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 103408). For *in vivo* studies or highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 103430) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/microg).

Application References: 1. Kato K, *et al.* 1992. *J. Exp. Med.* 176:1241. (IP, Costim, Block)
2. Qin L, *et al.* 1994. *J. Exp. Med.* 179:341. (Costim, Block)

Description: CD48 is a 45 kD GPI-anchored glycoprotein also known as BCM1, Blast-1 (human), and OX-45 (rat). It is a member of the Ig superfamily, expressed on T and B cells and monocytes/macrophages. It plays a role in adhesion and T cell recognition. The primary ligands for CD48 are CD2 and CD244.

* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.

- Antigen**
- References:**
1. Barclay AN, *et al.* 1997. The Leukocyte Antigen FactsBook Academic Press.
 2. Flament C, *et al.* 1996. *Hum. Immunol.* 46:82.
 3. Van der Merwe PA, *et al.* 1995. *Curr. Biol.* 5:74.
 4. Latchman Y