

PE/Cy7 anti-mouse CD49b

Catalog # / Size: 1117585 / 25 µg
1117590 / 100 µg

Clone: HMα2

Isotype: Hamster IgG


Immunogen: Mouse colon carcinoma cell line Colon26

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2



C57BL/6 splenocytes were stained with NK1.1 APC and CD49b (clone HMα2) PE/Cy7 (top), or Armenian Hamster IgG PE/Cy7 isotype control (bottom).

Applications:


Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.125 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation¹, immunofluorescence on frozen sections⁴, and blocking of cell adhesion¹⁻³. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays.

Application References:

1. Miyake S, *et al.* 1994. *Eur. J. Immunol.* 24:2000. (Block, IP)
2. Noto K, *et al.* 1995. *Int. Immunol.* 7:835. (Block)
3. Arase H, *et al.* 2001. *J. Immunol.* 167:1141. (Block)
4. Zhang Z, *et al.* 2008. *Blood* 111:1980. (IF)



Description: CD49b is a 150 kD glycoprotein, also known as α₂ integrin, VLA-2 α chain, Integrin α₂ chain, and HMα2. It is a member of the integrin family, expressed on NK cells, a subset of splenic CD4⁺ T cells, NK-T cells, intestinal intraepithelial and lamina propria lymphocytes, epithelial cells, and platelets. By associating with CD29 (integrin β₁ subunit), CD49b forms the VLA-2 (integrin α₂β₁) complex. It plays a

critical role in both adhesion and lymphocyte activation. The primary ligands for CD49b are collagen, laminin, and fibronectin. The HM α 2 antibody has been shown to be useful for partially blocking CD49b mediated interactions with collagen. Additionally, this antibody blocks staining of splenic NK cells by the monoclonal antibody DX5.

**Antigen
References:**

1. Barclay AN, *et al.* 1997. The Leukocyte Antigen FactsBook Academic Press.
2. Sasaki K, *et al.* 2003. *Int. Immunol.* 15:701.
3. Inoue O, *et al.* 2003. *J. Cell Biol.* 160:769.