


Purified anti-mouse/rat CD61

Catalog # / Size: 1121510 / 500 µg
Clone: 2C9.G2 (HMβ3-1)
Isotype: Hamster IgG
Immunogen: Vitronectin receptor protein from the mouse T-cell hybridoma 2B4
Reactivity: Mouse,Rat
Preparation: The antibody was purified by affinity chromatography.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration: 0.5



C57BL/6 mouse bone marrow cells stained with purified 2C9.G2, followed by anti-Armenian hamster IgG FITC

Applications:

Applications: Flow Cytometry, Immunohistochemistry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 1.0 microg per 10^6 cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: blocking of ligand binding¹⁻⁴, activation of $\alpha_v \beta_3$ integrin signaling⁵, and immunohistochemical staining of acetone-fixed frozen sections. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 104310).

Application References:

1. Kieffer N, *et al.* 1990. *Annu. Rev. Cell Biol.* 6:329. (Block)
2. Piali L, *et al.* 1995. *J. Cell Biol.* 130:451. (Block)
3. Ashkar S, *et al.* 2000. *Science* 287:860. (Block)
4. Schultz JF, *et al.* 1995. *J. Biol. Chem.* 270:11522. (Block)
5. Moulder K, *et al.* 1991. *J. Exp. Med.* 173:343. (Activ)
6. Carlson TR, *et al.* 2008.135:2193. [PubMed](#)
7. Yamaji D, *et al.* 2009. *Genes Dev.* 23:2382. [PubMed](#)

Description: CD61 is a 110 kD integrin β chain also known as β_3 integrin or gpIIb. It associates with the integrin α_v chain (CD51) to form the vitronectin receptor. In addition, CD61 can associate with the integrin α_{IIb} chain (CD41) to form the gpIIb/IIIa complex. CD61 is expressed on platelets, megakaryocytes, endothelium, smooth muscle, a subset of B cells, myeloid cells, osteoclasts, and mast cells. CD61, in conjunction with CD41 or CD51, mediates adhesion to fibronectin, fibrinogen, vitronectin, thrombospondin, and von Willebrand factor. Leukocyte-endothelial adhesion is mediated by the binding of α_v/β_3 integrin or vitronectin receptor to CD31 (PECAM-1).

Antigen 1. Barclay A, *et al.* 1997. The Leukocyte Antigen FactsBook. Academic Press.

- References:**
2. Phillips DR, *et al.* 1991. *Cell* 65:359.
 3. Felding-Habermann B, *et al.* 1993. *Curr. Opinion Cell Biol.* 5:864.