


**Biotin anti-mouse/rat CD61**

**Catalog # / Size:** 1121520 / 500 µg  
**Clone:** 2C9.G2 (HMβ3-1)  
**Isotype:** Hamster IgG  
**Immunogen:** Vitronectin receptor protein from the mouse T-cell hybridoma 2B4  
**Reactivity:** Mouse,Rat  
**Preparation:** The antibody was purified by affinity chromatography, and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin.  
**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.  
**Concentration:** 0.5



C57BL/6 mouse bone marrow cells stained with biotinylated 2C9.G2, followed by Sav-PE

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 microg per 10<sup>6</sup> cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

**Application Notes:** Additional reported applications (for the relevant formats) include: blocking of ligand binding<sup>1-4</sup>, activation of α<sub>v</sub> β<sub>3</sub> integrin signaling<sup>5</sup>, and immunohistochemical staining of acetone-fixed frozen sections. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 104310).

- Application References:**
1. Kieffer N, *et al.* 1990. *Annu. Rev. Cell Biol.* 6:329. (Block)
  2. Piali L, *et al.* 1995. *J. Cell Biol.* 130:451. (Block)
  3. Ashkar S, *et al.* 2000. *Science* 287:860. (Block)
  4. Schultz JF, *et al.* 1995. *J. Biol. Chem.* 270:11522. (Block)
  5. Moulder K, *et al.* 1991. *J. Exp. Med.* 173:343. (Activ)
  6. Carlson TR, *et al.* 2008.135:2193. [PubMed](#)
  7. Yamaji D, *et al.* 2009. *Genes Dev.* 23:2382. [PubMed](#)
  8. Milani ES, *et al.* 2013. *Development.* 140:117. [PubMed](#).

**Description:** CD61 is a 110 kD integrin β chain also known as β<sub>3</sub> integrin or gpIIIa. It associates with the integrin α<sub>v</sub> chain (CD51) to form the vitronectin receptor. In addition, CD61 can associate with the integrin α<sub>IIb</sub> chain (CD41) to form the gpIIb/IIIa complex. CD61 is expressed on platelets, megakaryocytes, endothelium, smooth muscle, a subset of B cells, myeloid cells, osteoclasts, and mast cells. CD61, in conjunction with CD41 or CD51, mediates adhesion to fibronectin, fibrinogen, vitronectin, thrombospondin, and von Willebrand factor. Leukocyte-endothelial adhesion is mediated by the binding of α<sub>v</sub>/β<sub>3</sub> integrin or vitronectin receptor to CD31 (PECAM-1).

**Antigen** 1. Barclay A, *et al.* 1997. The Leukocyte Antigen FactsBook Academic Press.

- References:**
2. Phillips DR, *et al.* 1991. *Cell* 65:359.
  3. Felding-Habermann B, *et al.* 1993. *Curr. Opinion Cell Biol.* 5:864.