

PE/Cy7 anti-mouse CD94

Catalog # / Size: 1127545 / 50 µg
1127550 / 200 µg

Clone: 18d3

Isotype: Rat IgG2a, κ


Immunogen: CHO cells expressing the B6 alleles of CD94 and NKG2A

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: Lot-specific




C57BL/6 mouse splenocytes were stained with CD49b (pan-NK cells) FITC and CD94 (clone 18d3) PE/Cy7 (top), or rat IgG2a, κ PE/Cy7 isotype control (bottom).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.06 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.



Application References: 1. Vance RE, et al. 1999. *J. Exp. Med.* 190:1801.
2. Tang X, et al. 2006. *J. Immunol.* 177:7645. [PubMed](#)

This product may be used for research purposes only. It is not licensed for resale and may only be used by the buyer. This product may not be used and is not licensed for clinical assays where the results of such assays are provided as a diagnostic service. If a diagnostic or therapeutic use is anticipated then a license must be requested from the University of California. The availability of such diagnostic and therapeutic use license(s) cannot be guaranteed from the University of California.

Description: CD94 is a 43/39 kD C-type lectin, also known as Kp43. It is present on all NK cells, NKT cells, and a subset of CD8-positive T lymphocytes in most mouse strains. CD94 is a type-II transmembrane protein with an extracellular lectin-like domain and a short cytoplasmic tail. CD94 is expressed as a disulphide-linked heterodimer with a NKG2 subunit believed to mediate signal transduction. When associated with NKG2A, the complex triggers inhibition; when associated with NKG2C, the complex triggers stimulation. The receptor complex of CD94 and NKG2 receptors bind to the ligand, Qa-1, and are thought to play a role in maintaining self-tolerance in developing NK cells.

- Antigen**
- References:**
1. Barclay A, *et al.* 1997. The Leukocyte Antigen FactsBook Academic Press.
 2. Lopez-Botet M, *et al.* 1997. *Immunol. Rev.* 155:165.
 3. Moretta A, *et al.* 1997. *Immunol. Rev.* 155:105.
 4. Phillip