SONY

Product Data Sheet

APC anti-mouse CD152

Catalog # / Size:	1131550 / 100 µg 1131545 / 25 µg
Clone:	UC10-4B9
Isotype:	Hamster IgG
Immunogen:	Mouse CTLA-4-mouse IgG2a fusion protein
Reactivity:	Mouse
Preparation:	The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Workshop Number:	750 under optimal conditions.
Concentration:	0.2



Con A+IL-2-stimulated C57BL/6 splenocytes (3 days) stained with UC10-4B9 APC and CD3e (145-2C11) PE

Applications:

Applications:	Flow Cytometry
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤1.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.



Con A+IL-2-stimulated C57BL/6 splenocytes (3 days) stained with Armenian Hamster IgG isotype APC and CD3e (145-2C11) PE

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.
Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com

Application Notes: The UC10-4B9 antibody can enhance T cell co-stimulation by blocking CTLA-4 interactions with the B7 co-receptors, favoring CD28 interactions. Additional reported applications (for the relevant formats) include: immunoprecipitation1, *in vitro* stimulation, *in vitro* and *in vivo* blocking[1-4] of ligand binding, and as Other capture antibody5. To reduce non-specific binding to cells bearing Fc-receptors, pre-incubation of cells with anti-mouse CD16/CD32, clone 93 (Cat. No. 101301/101302), is recommended prior to immunofluorescent staining. For most successful immunofluorescent staining results, it may be important to maximize signal over background by using a relatively bright fluorochrome-antibody conjugate (Cat. No. 106306) or by using a high sensitivity, three-layer staining technique (e.g., including a biotinylated anti-Armenian hamster IgG (Cat. No. 405501) second step, followed by SAv-PE (Cat. No. 405204)). The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 106308).

Application References:

1. Walunas TL, *et al.* 1994. *Immunity* 1:405. (Block, IP)
2. Cilio CM, *et al.* 1998. *J. Exp. Med.* 188:1239. (Block)
3. Issazadeh S, *et al.* 1999. *J. Immunol.* 162:761. (Block)
4. McCoy K, *et al.* 1997. *J. Exp. Med.* 186:183. (Block)
5. Hsu HC, *et al.* 2007. *J. Immunol.* 178:5357. (Other Capture)
6. Sugita S, *et al.* 2010. *Invest. Ophthalmol. Vis. Sci.* 51:5783. PubMed

---

Description: CD152, also known as CTLA-4 or Ly-56, is a 33 kD member of the immunoglobulin superfamily. It is expressed on activated T and B lymphocytes. CD152 is similar to CD28 in amino acid sequence, structure, and genomic organization and these two receptors share common B7 family counter-receptors (B7-1, B7-2). Whereas CD28 delivers a costimulatory signal in T cell activation, CTLA-4 negatively regulates cell-mediated immune responses. CD152 is thought to play a role in the induction and maintenance of immunological tolerance as well as the development of protective immunity and thymocyte regulation.

Antigen References:

1. Barclay A, *et al.* 1997. The Leukocyte Antigen FactsBook Academic Press.
2. Allison JP, *et al.* 1995. *Science* 270:932.
3. Waterhouse P, *et al.* 1995. *Science* 270:985.
4. Linsley PS, *et al.* 1991. *J. Exp. Med.* 174:561.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.