SONY

Product Data Sheet

PerCP/Cyanine5.5 anti-mouse CD154

Catalog # / Size: 1132570 / 100 µg
1132565 / 25 µg

Clone: MR1

Isotype: Hamster IgG

Immunogen: Activated mouse Th1 clone D1.6

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with PerCP/Cyanine5.5 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Workshop Number: 750 under optimal conditions.

Concentration: 0.2



Enriched C57BL/6 mouse splenic T cells were stimulated with PMA+ionomycin for 6 hours and then stained with CD154 (clone MR1) PerCP/Cyanine5.5 (filled histogram) or Armenian hamster IgG PerCP/Cyanine5.5 (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.5 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

* PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen sections[1,2], and *in vitro* and *in vivo* blocking of ligand binding[3-5]. For most successful immunofluorescent staining results, it may be important to maximize signal over background by using a relatively bright fluorochrome-antibody conjugate (Cat. No. 106506) or by using a high sensitivity, three-layer staining technique (e.g., including a biotinylated antibody (Cat. No. 106504) or biotinylated anti-Armenian hamster IgG (Cat. No. 405501) second step, followed by SAv-PE (Cat. No. 45204)). The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 106508).

Application References:

1. Lettesjö H, *et al.* 2000. *J. Immunol.* 165:4095. (IHC)
2. Dunn RJ, *et al.* 1997. *J. Histochem. Cytochem.* 45:129. (IHC)
3. Noelle RJ, *et al.* 1992. *P. Natl. Acad. Sci. USA* 89:6550. (Block)
4. Roy M, *et al.* 1995. *Eur. J. Immunol.* 25:596. (Block)
5. Foy TM, *et al.* 1994. *J. Exp. Med.* 180:157. (Block)
6. Lawson BR, *et al.* 2007. *J. Immunol.* 178:5366.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.

Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com

Description: CD154 is a 39 kD TNF superfamily member also known as CD40 ligand, gp39, T-BAM, TRAP, and Ly-62. CD154 is an accessory molecule expressed predominantly on activated $CD4^+$ lymphocytes that bind CD40. CD154 plays an important role in T-B cell costimulation. The MR1 antibody has been reported to inhibit the activation of T and B lymphocytes *in vitro* and antigen-specific lymphocyte responses *in vivo*.

Antigen References:

1. Barclay A, *et al.* 1997. The Leukocyte Antigen FactsBook Academic Press.
2. Noelle RJ, *et al.* 1992. *P. Natl. Acad. Sci. USA* 89:6550.
3. Bancherou J, *et al.* 1994. *Annu. Rev. Immunol.* 12:881.
4. Clark EA, *et al.* 1996. *P. Natl. Acad. Sci. USA* 83:4494.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.