

Brilliant Violet 421™ anti-mouse Ly-6A/E (Sca-1)

Catalog # / Size: 1140635 / 125 µl
1140640 / 50 µg

Clone: D7

Isotype: Rat IgG2a, κ


Immunogen: IL-2-dependent mouse T-cell line (CTL-L)

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 421™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 421™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Concentration: microg sizes: 0.2 mg/ml
microL sizes: lot-specific



C57BL/6 mouse splenocytes were stained with Ly-6A/E (clone D7) Brilliant Violet 421™ (filled histogram) or rat IgG2a, κ Brilliant Violet 421™ isotype control (open histogram).


Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining using the microg size, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. For immunofluorescent staining using the microL size, the suggested use of this reagent is ≤5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use



C57BL/6 mouse bone marrow cells were stained with Ly-6A/E (clone D7) Brilliant Violet 421™ (filled histogram) or rat IgG2a, κ Brilliant Violet 421™ isotype control (open histogram). Data shown was gated on lymphoid cell population.

for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: The D7 antibody has been reported to induce T cell activation and inhibit TCR-induced IL-2 production. Additional reported applications (for the relevant formats) include: Western blotting^{1,2}, immunoprecipitation¹, *in vitro* lymphocyte activation³⁻⁶, induction of redirected lysis⁷, induction of T cell inhibitory signalling⁸, immunofluorescence⁹, and immunohistochemical staining of acetone-fixed frozen sections¹³ and Bouin-fixed, paraffin-embedded samples⁹.

The two Sca-1 recognizing clones D7 and [E13-161.7](#) have been shown to bind distinct epitopes due to the inability of D7 to block the binding of E13-161.7.¹⁴

- Application References:**
1. Ortega G, *et al.* 1986. *J. Immunol.* 137:3240. (WB, IP)
 2. Palfree RGE, *et al.* 1986. *Immunogenetics* 23:197. (WB)
 3. Codias EK, *et al.* 1990. *J. Immunol.* 144:2197.
 4. Malek TR, *et al.* 1986. *J. Exp. Med.* 164:709.
 5. Codias EK, *et al.* 1990. *J. Immunol.* 145:1407.
 6. Ivanov V, *et al.* 1994. *J. Immunol.* 153:2394.
 7. Karlhofer FM, *et al.* 1991. *J. Immunol.* 146:3662.
 8. Fleming T, *et al.* 1994. *J. Immunol.* 153:1955.
 9. van Bragt MPA, *et al.* 2005. *Biol. Reprod.* 73:634. (IF, IHC)
 10. Umland O, *et al.* 2007. *J. Immunol.* 178:4147.
 11. Cridland SO, *et al.* 2009. *Blood Cell. Mol. Dis.* 45:149. (FC) [PubMed](#)
 12. Pronk CJ, *et al.* 2011. *J. Exp Med.* [PubMed](#)
 13. English A, *et al.* 2000. *J. Immunol.* 165:3763. (IHC)
 14. Bamezai A and Rock KL. 1995. *Proc. Natl. Acad. Sci. USA* 92:4294.

Description: Ly-6A/E, also known as Sca-1, is an 18 kD member of the Ly-6 multigene family. Ly6A/E is a glycosylphosphatidylinositol (GPI)-linked protein expressed on hematopoietic stem cells. In mice expressing the Ly-6.2 haplotype (e.g., AKR, C57BL, C57BR, DBA/2, SJL, SWR, and 129), Ly-6A/E is also expressed on peripheral B lymphocytes and thymic and peripheral T lymphocytes. Strains expressing the Ly-6.1 haplotype (e.g., BALB/c, CBA, C3H/He, DBA/1, and NZB) have low Ly-6A/E expression on resting peripheral lymphocytes. The expression of Ly-6A/E on lymphocytes is upregulated upon activation from both Ly6.1 and Ly6.2 haplotype mice. Ly-6A/E is thought to be involved in the regulation of both T and B cell responses.

- Antigen References:**
1. Rock KL, *et al.* 1989. *Immunol. Rev.* 111:195.
 2. Morrison SJ, *et al.* 1994. *Immunity* 1:661.
 3. Spangrude GJ, *et al.* 1988. *J. Immunol.* 141:3697.
 4. Malek T, *et al.* 1986. <