

Alexa Fluor® 647 anti-mouse Ly-51

Catalog # / Size: 1141555 / 25 µg
1141560 / 100 µg

Clone: 6C3

Isotype: Rat IgG2a, κ


Immunogen: C57BL/6 mouse Pre-B lymphoma cell line (L1-2) plus Abelson murine leukemia virus-specific cytotoxic T-cell clones

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with Alexa Fluor® 647 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5




C57BL/6 mouse bone marrow cells stained with CD45R (RA3-6B2) PE and Alexa Fluor® 647 rat IgG2a isotype control

Applications:

Applications: Immunofluorescence

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤1.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.



C57BL/6 mouse bone marrow cells stained with CD45R (RA3-6B2) PE and 6C3 Alexa Fluor® 647

* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633 nm / 635 nm.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation^{1,2} and immunohistochemical staining^{3-5, 7-8} of acetone-fixed frozen sections.

- Application References:**
1. Pillemer E, *et al.* 1984. *P. Natl. Acad. Sci. USA* 81:4434. (IP)
 2. Hardy RR, *et al.* 1991. *J. Exp. Med.* 173:1213. (IP)
 3. Adkins B, *et al.* 1988. *Immunogenetics* 27:180. (IHC)
 4. Surh CD, *et al.* 1992. *J. Exp. Med.* 176:611. (IHC)
 5. Goverman J, *et al.* 1997. *Immunol. Today* 18:204. (IHC)
 6. Dumont-Lagace M, 2014. *J. Immunol.* 192:2219. [PubMed](#)
 7. Kvell K, *et al.* 2010. *PLoS One* 5:e10701. (FC, IHC)
 8. Griewank K, *et al.* 2007. *Immunity* 27:751. (FC, IHC)

Description: Ly-51 is a 140 kD protein also known as 6C3/BP-1. It is a homodimeric cell-surface glycoprotein with aminopeptidase A (APA) activity. Ly-51 is expressed on B cell

progenitors, bone marrow stromal cell lines, thymic dendritic cells, and cortical epithelial cells. Ly-51 expression can be upregulated by IL-7 stimulation.

**Antigen
References:**

1. Goverman J, *et al.* 1997. *Immunol. Today* 18:204.
2. Morse HC, *et al.* 1987. *J. Exp. Med.* 165:920.
3. Sherwood P, *et al.* 1990. *Int. Immunol.* 2:399.
4. Lin Q, *et al.* 1998.