Product Data Sheet

Purified anti-mouse CD279 (PD-1)

Catalog # / 1145505 / 50 µg

Size:

Clone: RMP1-30

Isotype: Rat IgG2b, ĸ

Mouse PD-1 transfected BHK cells Immunogen:

Reactivity: Mouse


The antibody was purified by affinity **Preparation:**

chromatography.

Phosphate-buffered solution, pH 7.2, Formulation:

containing 0.09% sodium azide.

Concentration: 0.5



Con A (3-day) activated C57BL/6 mouse splenocytes stained with RMP1-30 PE

Applications:

Applications: Other

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL

volume. It is recommended that the reagent be titrated for optimal

performance for each application.

Application Notes: Additional reported application (for the relevant formats) include: Functional assay. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 109108). The RMP1-30 antibody does not block the binding of PD-1 to B7-H1 and B7-DC1.

Application

1. Matsumoto K, et al. 2004. J. Immunol. 172:2530.

References:

2. Raimondi G, et al. 2006. J. Immunol. 176:2808. (FC) PubMed

3. King IL, et al. 2009. J. Exp Med 206:1001. (FC) PubMed

Description:

CD279 is a 50-55 kD immunoglobulin superfamily member also known as programmed death-1 (PD-1). PD-1 is expressed on a subset of CD4⁻CD8⁻ thymocytes and on activated T and B cells. PD-1 is thought to be involved in lymphocyte clonal selection and peripheral tolerance. The PD-1 ligands, PD-L1 (also known as B7-H1) and PD-L2 (B7-DC), are members of the B7 immunoglobulin superfamily.

Antigen References: 1. Barclay A, et al. 1997. The Leukocyte Antigen FactsBook Academic Press.

2. Agata Y. et al. 1996. Int. Immunol. 8:765.

3. Nishimura H, et al. 2001. Science 291:319.

4. Ishida Y, et al