## Alexa Fluor® 647 anti-mouse CD279 (PD-1)

Catalog # / Size: 1145585 / 25 µg

1145590 / 100 µg

Clone: RMP1-30 Isotype: Rat IgG2b, ĸ

Mouse PD-1 transfected BHK cells Immunogen:

Reactivity: Mouse

**Preparation:** The antibody was purified by affinity

chromatography and conjugated with


Alexa Fluor® 647 under optimal

conditions.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

**Concentration:** 0.2



C57BL/6 splenocytes were stained with CD3 PE and CD279 (clone RMP1-30) Alexa Fluor® 647 (top) or rat IgG2b, κ Alexa Fluor® 647 isotype control (bottom).

CD3 PE

105

105

104

103

Rat IgG2b, x Isotype Control

## **Applications:**

**Applications:** Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤1.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

\* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at

633 nm / 635 nm.

**Application** Notes: Additional reported application (for the relevant formats) include: Functional assay. The LEAF  $^{\text{\tiny{TM}}}$  purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 109108). The RMP1-30 antibody does not block the binding of PD-1 to B7-H1 and B7-DC1.


1. Matsumoto K, et al. 2004. J. Immunol. 172:2530.

**Application** References: 2. Raimondi G, et al. 2006. J. Immunol. 176:2808. (FC) PubMed

3. King IL, et al. 2009. J. Exp Med 206:1001. (FC) PubMed

**Description:** CD279 is a 50-55 kD immunoglobulin superfamily member also known as

programmed death-1 (PD-1). PD-1 is expressed on a subset of CD4<sup>-</sup>CD8<sup>-</sup>



thymocytes and on activated T and B cells. PD-1 is thought to be involved in lymphocyte clonal selection and peripheral tolerance. The PD-1 ligands, PD-L1 (also known as B7-H1) and PD-L2 (B7-DC), are members of the B7 immunoglobulin superfamily.

## Antigen References:

- 1. Barclay A, et al. 1997. The Leukocyte Antigen FactsBook Academic Press.
- 2. Agata Y, et al. 1996. Int. Immunol. 8:765.
- 3. Nishimura H, et al. 2001. Science 291:319.
- 4. Ishida Y, et al