

**PE/Cy7 anti-mouse TCR β chain**

**Catalog # / Size:** 1146110 / 100 μg  
1146105 / 25 μg

**Clone:** H57-597

**Isotype:** Hamster IgG


**Immunogen:** Affinity purified TCR from mouse DO-11.10 cells

**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.2



C57BL/6 mouse splenocytes stained with H57-597 PE/Cy7

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 microg per 10<sup>6</sup> cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

**Application Notes:** H57-597 is a hamster mAb directed to an epitope of the C region of TCR β chain<sup>12</sup>. The H57-597 antibody does not cross-react with γ/δ TCR-bearing T cells. Immobilized or soluble H57-597 antibody can activate α/β TCR-bearing T cells. Additional reported applications (for the relevant formats) for this antibody include: immunoprecipitation<sup>2</sup>, *in vitro* stimulation<sup>2,3</sup>, *in vivo* depletion<sup>4-6</sup>, and immunohistochemical staining of acetone-fixed frozen sections<sup>7,8,9</sup>. The LEAF™ purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 109214).

**Application References:**

1. Gascoigne NJ. 1990. *J. Biol. Chem.* 265:9296.
2. Kruisbeek A, *et al.* 1991. *In Current Protocols in Immunology.* pp. 3.12.1. (Costim IP)
3. Davenport C, *et al.* 1995. *J. Immunol.* 155:3742. (Costim)
4. Drobyski W, *et al.* 1996. *Blood* 87:5355. (Deplete)
5. Kummer U, *et al.* 2001. *Immunol. Lett.* 75:153. (Deplete)
6. van der Heyde HC, *et al.* 1995. *J. Immunol.* 154:3985. (Deplete)
7. Tomita K, *et al.* 1999. *Genes Dev.* 13:1203. (IHC)
8. Podd BS, *et al.* 2006. *J. Immunol.* 176:6532. (IHC)
9. Ponomarev ED, *et al.* 2007. *J. Immunol.* 178:39. (IHC)
10. Chappaz S, *et al.* 2007. *Blood* doi:10.1182/blood-2007-02-074245. (FC)
11. Tsukumo S, *et al.* 2006. *J. Immunol.* 177:8365. (FC) [PubMed](#)
12. Grégoire C, *et al.* 1991. *Proc. Natl. Acad. Sci USA* 88:8077.

**Description:** T cell receptor (TCR) is a heterodimer consisting of an α and a β chain (TCR α/β)

or a  $\gamma$  and a  $\delta$  chain (TCR  $\gamma/\delta$ ). TCR- $\beta$  is a member of the immunoglobulin superfamily and a component of the CD3/TCR complex (along with TCR- $\alpha$ ). It is expressed on  $\alpha/\beta$  TCR-bearing T cells and thymocytes. The CD3/TCR complex plays a key role in antigen recognition, signal transduction, and T cell activation.

**Antigen  
References:**

1. Davis MM, *et al.* 1998. *Ann. Rev. Immunol.* 16:523.
2. Huppa JB, *et al.* 2003. *Nat. Immunol.* 4:749.
3. Kubo R, *et al.* 1989. *J. Immunol.* 142:2736.