

FITC anti-mouse H-2D b

Catalog # / Size: 1157530 / 500 µg
1157525 / 50 µg

Clone: KH95

Isotype: Mouse IgG2b, κ


Immunogen: C57BL/10 mouse skin graft and splenocytes

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5




BALB/C mouse splenocytes were stained with H-2b (clone KH95) FITC (filled histogram), or mouse IgG2b, κ FITC isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 1.0 microg per 10⁶ cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: complement-dependent cytotoxicity¹, and Western blotting.



C57BL/6 mouse splenocytes were stained with H-2b (clone KH95) FITC (filled histogram) or mouse IgG2b, κ FITC isotype control (open histogram).

- Application References:**
1. Hasenkrug KJ, *et al.* 1987. *Immunogenetics* 25:136.
 2. Shao H, *et al.* 2005. *J. Immunol.* 175:1851.
 3. Ponomarev ED, *et al.* 2006. *J. Immunol.* 176:1402.
 4. Robb RJ, *et al.* 2012 *Blood.* 119:5898. [PubMed](#)
 5. Zhang P, *et al.* 2013. *J. Immunol.* 191:5291. [PubMed](#)
 6. Quinn KM, *et al.* 2013. *J. Immunol.* 191:5085. [PubMed](#)
 7. Markey KA, *et al.* 2014. *J Immunol.* 192:5426. [PubMed](#)
 8. Hogan T, *et al.* 2014. *PLoS Comput Biol.* 10:1003805. [PubMed](#)

Description: The KH95 antibody reacts with the H-2Db MHC class I alloantigen expressed on nucleated cells from mice of the H-2Db haplotype. H-2Db is involved in antigen presentation to T cells expressing CD3/TCR and CD8 proteins. Reactivity with other haplotypes (*e.g.*, a,d,f,k,n,p,q,r,s,u,v) has not been reported.

- Antigen**
- References:**
1. Watts C. 1997. *Ann. Rev. Immunol.* 15:821.
 2. Pamer E, *et al.* 1998. *Ann. Rev. Immunol.* 16:323.
 3. York I, *et al.* 1996. *Ann. Rev. Immunol.* 14:369.