Product Data Sheet

APC anti-mouse H-2D b

Catalog # / Size: 1157565 / 25 µg

1157570 / 100 µg

Clone:

Isotype: Mouse IgG2b, κ

C57BL/10 mouse skin graft and Immunogen:

splenocytes

Reactivity: Mouse

Preparation: The antibody was purified by affinity


> chromatography and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and

unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

Concentration: 0.2



C57BL/6 mouse splenocytes were stained with H-2b (clone KH95) APC (filled histogram) or mouse IgG2b, K

APC isotype control (open

histogram).

Applications:

Applications: Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of

this reagent is ≤0.5 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each

application.

Application Notes:

Additional reported applications (for the relevant formats) include: complement-

dependent cytotoxicity1, and Western blotting.

Application References: 1. Hasenkrug KJ, et al. 1987. Immunogenetics 25:136.

2. Shao H, et al. 2005. J. Immunol. 175:1851.

3. Ponomarev ED, et al. 2006. J. Immunol. 176:1402. 4. Robb RJ, et al. 2012 Blood. 119:5898. PubMed

5. Zhang P, et al. 2013. J. Immunol. 191:5291. PubMed 6. Quinn KM, et al. 2013. J. Immunol. 191:5085. PubMed

7. Markey KA, et al. 2014. J Immunol. 192:5426. PubMed

8. Hogan T, et al. 2014. PLoS Comput Biol. 10:1003805. PubMed

Description: The KH95 antibody reacts with the H-2Db MHC class I alloantigen expressed on

nucleated cells from mice of the H-2Db haplotype. H-2Db is involved in antigen presentation to T cells expressing CD3/TCR and CD8 proteins. Reactivity with

other haplotypes (e.g., a,d,f,k,n,p,q,r,s,u,v) has not been reported.

Antigen References: 1. Watts C. 1997. Ann. Rev. Immunol. 15:821.

2. Pamer E, et al. 1998. Ann. Rev. Immunol. 16:323.

3. York I, et al. 1996. Ann. Rev. Immunol. 14:369.