Product Data Sheet

PE/Dazzle™ 594 anti-mouse H-2D b

Catalog # / 1157610 / 100 µg

Size: 1157605 / 25 µg

Clone: **KH95**

Isotype: Mouse IgG2b, κ

C57BL/10 mouse skin graft and Immunogen:

splenocytes

Reactivity: Mouse

The antibody was purified by affinity Preparation:


> chromatography and conjugated with PE/Dazzleâ,,¢ 594 under optimal conditions. The solution is free of unconjugated PE/Dazzleâ,,¢ 594 and

unconjugated antibody.

Phosphate-buffered solution, pH 7.2, Formulation:

containing 0.09% sodium azide.

Concentration: 0.2 ma/ml



C57BL/6 mouse splenocytes were stained with H-2Db (clone KH95) PE/Dazzle™ 594 (filled histogram) or Mouse IgG2b, κ PE/Dazzle[™] 594 isotype control (open histogram).

Applications:

Flow Cytometry **Applications:**

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.5 μg per million cells in 100 μl volume. It is recommended that the reagent be titrated for optimal performance for each application.

* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum

emission of 610 nm.

Application Notes:

Additional reported applications (for the relevant formats) include:

complement-dependent cytotoxicity¹, and Western blotting.

Application

1. Watts C. 1997. Ann. Rev. Immunol. 15:821.

References:

2. Pamer E, et al. 1998. Ann. Rev. Immunol. 16:323.

3. York I. et al. 1996. Ann. Rev. Immunol. 14:369.

Description: The KH95 antibody reacts with the H-2Db MHC class I alloantigen expressed

on nucleated cells from mice of the H-2D^b haplotype. H-2D^b is involved in antigen presentation to T cells expressing CD3/TCR and CD8 proteins. Reactivity with other haplotypes (e.g., a,d,f,k,n,p,q,r,s,u,v) has not been

reported.

Antigen

1. Watts C. 1997. Ann. Rev. Immunol. 15:821.

References: 2. Pamer E, et al. 1998. Ann. Rev. Immunol. 16:323.

3. York I, et al. 1996. Ann. Rev. Immunol. 14:369.