


# APC/Fire™ 750 anti-mouse CD19

<b>Catalog # /</b>	1177785 / 25 µg
<b>Size:</b>	1177790 / 100 µg
<b>Clone:</b>	6D5
<b>Isotype:</b>	Rat IgG2a, κ
<b>Immunogen:</b>	Mouse CD19-expressing K562 human erythroleukemia cells
<b>Reactivity:</b>	Mouse
<b>Preparation:</b>	The antibody was purified by affinity chromatography and conjugated with APC/Fire™
<b>Formulation:</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
<b>Workshop Number:</b>	750 under optimal conditions.
<b>Concentration:</b>	0.2 mg/ml



C57BL/6 splenocytes were stained with CD3 PE and CD19 (clone 6D5) APC/Fire™ 750 (top) or Rat IgG2a, κ APC/Fire™ 750 isotype control (bottom).

## Applications:

**Applications:** Flow Cytometry


**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

\* APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 787 nm.

**Application Notes:** Additional reported applications (for the relevant formats) include: immunofluorescence<sup>7</sup>.

### Application References:

- Shoham T, et al. 2003. *J. Immunol.* 171:4062. (FC)
- Goodyear CS, et al. 2004. *J. Immunol.* 172:2870. (FC)
- Kamimura D, et al. 2006. *J. Immunol.* 177:306. (FC)
- Andoniou CE, et al. 2005. *Nat. Immunol.* 6:1011. (FC)
- Lawson BR, et al. 2007. *J. Immunol.* 178:5366. (FC)
- Phan TG, et al. 2007. *Nat. Immunol.* 8:992. (FC)
- Hayashida K, et al. 2008. *J. Biol. Chem.* 283:19895. (IF) [PubMed](#)
- Charles N, et al. 2010. *Nat. Med.* 16:701. (FC) [PubMed](#)
- Bankoti J, et al. 2010. *Toxicol. Sci.* 115:422. (FC) [PubMed](#)
- Stadnisky MD, et al. 2011. *Blood.* 117:5133. (FC) [PubMed](#)
- Perlot T, et al. 2012. *J. Immunol.* 188:1201. (FC) [PubMed](#)
- Olive V, et al. 2013. *Elife.* 2:822. [PubMed](#)
- Miyai T, et al. 2014. *PNAS.* 111:11780. [PubMed](#)



**Description:** CD19 is a 95 kD glycoprotein also known as B4. It is a member of the Ig superfamily, expressed on all pro-B to mature B cells (during development) and follicular dendritic cells. Plasma cells do not express CD19. CD19, in association with CD21 and CD81, forms a molecular complex integral to B cell activation.

**Antigen**  
**References:**

1. Fearon DT. 1993. *Curr. Opin. Immunol.* 5:341.
2. Krop I, et al. 1996. *Eur. J. Immunol.* 26:238.
3. Krop I, et al. 1996. *J. Immunol.* 157:48.
4. Tedder TF, et al. 1994. *Immunol. Today* 15:437.