

Spark Blue™ 550 anti-mouse CD19

Catalog # / Size: 1177830 / 100 µg
1177825 / 25 µg

Clone: 6D5

Isotype: Rat IgG2a, κ

Immunogen: Mouse CD19-expressing K562 human erythroleukemia cells


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with Spark Blue™ 550 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Workshop Number: 750 under optimal conditions.

Concentration: 0.5 mg/mL




C57BL/6 mouse splenocytes were stained with CD19 (clone 6D5) Spark Blue™ 550 (filled histogram.) Open histogram represents unstained cells.

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.5 µg per million cells in 100 µL volume. It is recommended that the reagent be titrated for optimal performance for each application.



* Spark Blue™ 550 has a maximum excitation of 516 nm and a maximum emission of 540 nm.

Application Notes: Additional reported applications (for the relevant formats) include: immunofluorescence⁷.

- Application References:**
1. Shoham T, et al. 2003. *J. Immunol.* 171:4062. (FC)
 2. Goodyear CS, et al. 2004. *J. Immunol.* 172:2870. (FC)
 3. Kamimura D, et al. 2006. *J. Immunol.* 177:306. (FC)
 4. Andoniou CE, et al. 2005. *Nat. Immunol.* 6:1011. (FC)
 5. Lawson BR, et al. 2007. *J. Immunol.* 178:5366. (FC)
 6. Phan TG, et al. 2007. *Nat. Immunol.* 8:992. (FC)
 7. Hayashida K, et al. 2008. *J. Biol. Chem.* 283:19895. (IF) [PubMed](#)
 8. Charles N, et al. 2010. *Nat. Med.* 16:701. (FC) [PubMed](#)
 9. Bankoti J, et al. 2010. *Toxicol. Sci.* 115:422. (FC) [PubMed](#)
 10. Stadnisky MD, et al. 2011. *Blood.* 117:5133. (FC) [PubMed](#)
 11. Perlot T, et al. 2012. *J. Immunol.* 188:1201. (FC) [PubMed](#)
 12. Olive V, et al. 2013. *Elife.* 2:822. [PubMed](#)
 13. Miyai T, et al. 2014. *PNAS.* 111:11780. [PubMed](#)

Description: CD19 is a 95 kD glycoprotein also known as B4. It is a member of the Ig superfamily, expressed on all pro-B to mature B cells (during development) and follicular dendritic cells. Plasma cells do not express CD19. CD19, in association with CD21 and CD81, forms a molecular complex integral to B cell activation.

Antigen
References:

1. Fearon DT. 1993. *Curr. Opin. Immunol.* 5:341.
2. Krop I, et al. 1996. *Eur. J. Immunol.* 26:238.
3. Krop I, et al. 1996. *J. Immunol.* 157:48.
4. Tedder TF, et al. 1994. *Immunol. Today* 15:437.