## PE/Cy7 anti-mouse TER-119/Erythroid Cells

Catalog # / Size: 1181110 / 100 μg

1181105 / 25 μg

Clone: TER-119

**Isotype:** Rat IgG2b, κ

**Immunogen:** Day-14 fetal liver cells from a C57BL/6

mouse

Reactivity: Mouse

**Preparation:** The antibody was purified by affinity


chromatography, and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7

and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

Concentration: 0.2



C57BL/6 mouse bone marrow cells stained with TER119 PE/Cy7

## **Applications:**

**Applications:** Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of

this reagent is  $\leq 0.25$  microg per 106 cells in 100 microL volume. It is

recommended that the reagent be titrated for optimal performance for each

application.

**Application** 

Notes:

The TER-119 antibody is useful for distinguishing erythrocytes and cells in the erythroid lineage. Additional reported applications (for the relevant formats) include: immunoprecipitation1, Western blotting1, complement-mediated cytotoxicity3, and immunohistochemical staining of acetone-fixed frozen sections and formalin-fixed paraffin-embedded sections. LEAF™ purified antibody

(Endotoxin < 0.1 EU/ $\mu$ g, Azide-Free, 0.2  $\mu$ m filtered) is recommended for

functional assays (Cat. No. 116214).

Application References:

1. Kina T, et al. 2000. Br. J. Haematol. 109:280. (IP, WB)

2. Vannucchi AM, et al. 2000. Blood 95:2559.

3. Maraskovsky E, et al. 1996. J. Exp. Med. 184:1953. (CMCD)

4. Grisendi S, *et al.* 2005. *Nature* 437:147. (FC)

5. Bourdeau A, et al. 2007. Blood 109:4220.

6. Chappaz S, et al. 2007. Blood 110:3862. (FC)

7. Heuser M, et al. 2007. Blood 110:1639. (FC)

**Description:** The TER-119 antigen is a 52 kD glycophorin A-associated protein, also known as

Ly-76. TER-119 is an erythroid-specific antigen expressed on early

proerythroblasts to mature erythrocytes, but not on erythroid colony-forming cells (BFU-E, blast-forming unit erythroid, or CFU-E, colony-forming unit erythroid).

Antigen References: 1. Kina T, et al. 2000. Br. J. Haematol. 109:280.

2. Ikuta K, *et al.* 1990. *Cell* 62:863.

3. Osawa M, et al. 1996. Weir's Handbook of Experimental Immunology. Vol. 2 pp.

66.1-66.5.