

**Pacific Blue™ anti-mouse TER-119/Erythroid Cells**

**Catalog # / Size:** 1181155 / 25 µg  
1181160 / 100 µg

**Clone:** TER-119

**Isotype:** Rat IgG2b, κ


**Immunogen:** Day-14 fetal liver cells from a C57BL/6 mouse

**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with Pacific Blue™ under optimal conditions. The solution is free of unconjugated Pacific Blue™.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.5



C57BL/6 bone marrow cells stained with Ter-119 Pacific Blue™

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 microg per 10<sup>6</sup> cells in 100 microL volume or 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

\* Pacific Blue™ has a maximum emission of 455 nm when it is excited at 405 nm. Prior to using Pacific Blue™ conjugate for flow cytometric analysis, please verify your flow cytometer's capability of exciting and detecting the fluorochrome.

**Application Notes:** The TER-119 antibody is useful for distinguishing erythrocytes and cells in the erythroid lineage. Additional reported applications (for the relevant formats) include: immunoprecipitation<sup>1</sup>, Western blotting<sup>1</sup>, complement-mediated cytotoxicity<sup>3</sup>, and immunohistochemical staining of acetone-fixed frozen sections and formalin-fixed paraffin-embedded sections. LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 116214).

- Application References:**
1. Kina T, *et al.* 2000. *Br. J. Haematol.* 109:280. (IP, WB)
  2. Vannucchi AM, *et al.* 2000. *Blood* 95:2559.
  3. Maraskovsky E, *et al.* 1996. *J. Exp. Med.* 184:1953. (CMCD)
  4. Grisendi S, *et al.* 2005. *Nature* 437:147. (FC)
  5. Bourdeau A, *et al.* 2007. *Blood* 109:4220.
  6. Chappaz S, *et al.* 2007. *Blood* 110:3862. (FC)
  7. Heuser M, *et al.* 2007. *Blood* 110:1639. (FC)

**Description:** The TER-119 antigen is a 52 kD glycophorin A-associated protein, also known as Ly-76. TER-119 is an erythroid-specific antigen expressed on early proerythroblasts to mature erythrocytes, but not on erythroid colony-forming cells (BFU-E, blast-forming unit erythroid, or CFU-E, colony-forming unit erythroid).

**Antigen** 1. Kina T, *et al.* 2000. *Br. J. Haematol.* 109:280.

- References:**
2. Ikuta K, *et al.* 1990. *Cell* 62:863.
  3. Osawa M, *et al.* 1996. *Weir's Handbook of Experimental Immunology*. Vol. 2 pp. 66.1-66.5.