


Brilliant Violet 510™ anti-mouse TER-119/Erythroid Cells


| | |
|--------------------------|--|
| Catalog # / Size: | 1181185 / 125 µl |
| Clone: | TER-119 |
| Isotype: | Rat IgG2b, κ |
| Immunogen: | Day-14 fetal liver cells from a C57BL/6 mouse |
| Reactivity: | Mouse |
| Preparation: | The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 510™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 510™ and unconjugated antibody. |
| Formulation: | Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA). |
| Concentration: | Lot-specific |



C57BL/6 bone marrow cells were stained with CD45 Pacific Blue™, and TER-119 Brilliant Violet 510™ (top) or rat IgG2b Brilliant Violet 510™ isotype control (bottom).

Applications:

| | |
|---------------------------|---|
| Applications: | Flow Cytometry |
| Recommended Usage: | Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. |



Brilliant Violet 510™ excites at 405 nm and emits at 510 nm. The bandpass filter 510/50 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel. Refer to your instrument manual or manufacturer for support. Brilliant Violet 510™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner.

into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: The TER-119 antibody is useful for distinguishing erythrocytes and cells in the erythroid lineage. Additional reported applications (for the relevant formats) include: immunoprecipitation¹, Western blotting¹, complement-mediated cytotoxicity³, and immunohistochemical staining of acetone-fixed frozen sections and formalin-fixed paraffin-embedded sections. LEAF™ purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 116214).

Application References:

1. Kina T, *et al.* 2000. *Br. J. Haematol.* 109:280. (IP, WB)
2. Vannucchi AM, *et al.* 2000. *Blood* 95:2559.
3. Maraskovsky E, *et al.* 1996. *J. Exp. Med.* 184:1953. (CMCD)
4. Grisendi S, *et al.* 2005. *Nature* 437:147. (FC)
5. Bourdeau A, *et al.* 2007. *Blood* 109:4220.
6. Chappaz S, *et al.* 2007. *Blood* 110:3862. (FC)
7. Heuser M, *et al.* 2007. *Blood* 110:1639. (FC)

Description: The TER-119 antigen is a 52 kD glycoporphin A-associated protein, also known as Ly-76. TER-119 is an erythroid-specific antigen expressed on early proerythroblasts to mature erythrocytes, but not on erythroid colony-forming cells (BFU-E, blast-forming unit erythroid, or CFU-E, colony-forming unit erythroid).

Antigen References:

1. Kina T, *et al.* 2000. *Br. J. Haematol.* 109:280.
2. Ikuta K, *et al.* 1990. *Cell* 62:863.
3. Osawa M, *et al.* 1996. *Weir's Handbook of Experimental Immunology*. Vol. 2 pp. 66.1-66.5.