

**APC/Fire™ 810 anti-mouse TER-119/Erythroid Cells**

**Catalog # / Size:** 1181320 / 100 µg  
1181315 / 25 µg

**Clone:** TER-119

**Isotype:** Rat IgG2b, κ

**Immunogen:** Day-14 fetal liver cells from a C57BL/6 mouse


**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 810 under optimal conditions.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide

**Workshop Number:** 750 under optimal conditions.

**Concentration:** 0.2 mg/mL




C57BL/6 bone marrow were stained with CD45 PE and TER-119 (clone TER-119) APC/Fire™ 810 (left) or CD45 PE only (right).

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 1.0 µg per million cells in 100 µL volume. It is recommended that the reagent be titrated for optimal performance for each application.



C57BL/6 mouse bone marrow cells were stained with CD150 (SLAM) (clone TC15-12F12.2) APC/Fire™ 750 (filled histogram) or rat IgG2a, κ APC/Fire™ 750 isotype control (open histogram).

\* APC/Fire™ 810 has a maximum excitation of 650 nm and a maximum emission of 810 nm.

**Application Notes:** The TER-119 antibody is useful for distinguishing erythrocytes and cells in the erythroid lineage. Additional reported applications (for the relevant formats) include: immunoprecipitation<sup>1</sup>, Western blotting<sup>1</sup>, complement-mediated cytotoxicity<sup>3</sup>, and immunohistochemical staining of acetone-fixed frozen sections and formalin-fixed paraffin-embedded sections. Ultra-LEAF™ purified antibody (Endotoxin < 0.01 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 116253-116258).

**Application  
References:**

1. Kina T, et al. 2000. *Br. J. Haematol.* 109:280. (IP, WB)
  2. Vannucchi AM, et al. 2000. *Blood* 95:2559.
  3. Maraskovsky E, et al. 1996. *J. Exp. Med.* 184:1953. (CMCD)
  4. Grisendi S, et al. 2005. *Nature* 437:147. (FC)
  5. Bourdeau A, et al. 2007. *Blood* 109:4220.
  6. Chappaz S, et al. 2007. *Blood* 110:3862. (FC)
  7. Heuser M, et al. 2007. *Blood* 110:1639. (FC)
  8. Gough SM, et al. 2014. *Cancer Discov.* 4:564. [PubMed](#)
- 

**Description:** The TER-119 antigen is a 52 kD glycoporphin A-associated protein, also known as Ly-76. TER-119 is an erythroid-specific antigen expressed on early proerythroblasts to mature erythrocytes, but not on erythroid colony-forming cells (BFU-E, blast-forming unit erythroid, or CFU-E, colony-forming unit erythroid).

**Antigen  
References:**

1. Kina T, et al. 2000. *Br. J. Haematol.* 109:280.
2. Ikuta K, et al. 1990. *Cell* 62:863.
3. Osawa M, et al. 1996. *Weir's Handbook of Experimental Immunology*. Vol. 2 pp. 66.1-66.5.