

FITC anti-mouse I-Ab

Catalog # / Size: 1182025 / 50 µg
1182030 / 500 µg

Clone: AF6-120.1

Isotype: Mouse IgG2a, κ


Immunogen: C57BL/10J splenocytes

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5



C57BL/6 mouse splenocytes stained with AF6-120.1 FITC

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤1.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for relevant formats of this clone) include: immunohistochemical staining of frozen sections (acetone-fixed⁵; OCT-embedded, ethanol-fixed sections⁷), immunofluorescence microscopy³ (including acetone-fixed epidermal sheets⁶), immunoprecipitation^{7,8}. Directly conjugated antibody was used for IF in (3) and (6) and for IHC in (5).

Application References:

1. Wall KA, *et al.* 1983. *J. Immunol.* 131:1056. (FC)
2. Cohn LE, *et al.* 1986. *P. Natl. Acad. Sci. USA* 83:747. (FC)
3. Inaba K, *et al.* 1998. *J. Exp. Med.* 188:2163 (IF)
4. Hamrah P, *et al.* 2002. *Invest Ophthalmol Vis. Sci.* 43:639 (IF)
5. Buono C, *et al.* 2003. *Arterioscler. Thromb. Vasc. Biol.* 23:454. (IHC)
6. Wang Z, *et al.* 2004. *J. Immunol.* 172:5924. (IHC IF)
7. Nakagawa TY, *et al.* 1999. *Immunity* 10:207. (IP)
8. Podolin PL, *et al.* 2008. *J. Immunol.* 180:7989. (FC IP) [PubMed](#)
9. Huang SC, *et al.* 2011. *J. Immunol.* 186:931. [PubMed](#)
10. Burghardt S, *et al.* 2013. *J. Immunol.* 191:5574. [PubMed](#)
11. Skrnjug I, *et al.* 2014. *PLoS One.* 9:110150. [PubMed](#)

Description: The AF6-120.1 antibody reacts with the I-Ab MHC class II alloantigen. These class II molecules are expressed on antigen presenting cells (including B cells) and a subset of T cells from H-2b bearing mice and are involved in antigen presentation to T cells expressing CD3/TCR and CD4 proteins. The AF6-120.1 antibody cross-reacts with H-2k and H-2^u haplotypes; this antibody does not cross-react with other haplotypes (d, f, q, r, s).

Antigen References:

1. Watts C. 1997. *Ann. Rev. Immunol.* 15:821.
2. Pamer E, *et al.* 1998. *Ann. Rev. Immunol.* 16:323.