PE anti-mouse I-Ab

Catalog # / Size: $1182035 / 50 \mu g$

1182040 / 200 µg

Clone: AF6-120.1

Isotype: Mouse IgG2a, κ

Immunogen: C57BL/10J splenocytes

Reactivity: Mouse

Preparation: The antibody was purified by affinity


chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and

unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

Concentration: 0.2



C57BL/6 mouse splenocytes stained with AF6-120.1 PE

Applications:

Applications: Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 microg per 10^6 cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application

Notes:

Additional reported applications (for relevant formats of this clone) include: immunohistochemical staining of frozen sections (acetone-fixed5; OCT-

embedded, ethanol-fixed sections⁷), immunofluorescence microscopy3 (including acetone-fixed epidermal sheets⁶), immunoprecipitation^{7,8}. Directly conjugated

antibody was used for IF in (3) and (6) and for IHC in (5).

Application References:

- 1. Wall KA, et al. 1983. J. Immunol. 131:1056. (FC)
- 2. Cohn LE, et al. 1986. P. Natl. Acad. Sci. USA 83:747. (FC)
- 3. Inaba K, et al. 1998. J. Exp. Med. 188:2163 (IF)
- 4. Hamrah P, et al. 2002. Invest Opthalmol Vis. Sci. 43:639 (IF)
- 5. Buono C, et al. 2003. Arterioscler. Thromb. Vasc. Biol. 23:454. (IHC)
- 6. Wang Z, et al. 2004. J. Immunol. 172:5924. (IHC IF) 7. Nakagawa TY, et al. 1999. Immunity 10:207. (IP)
- 8. Podolin PL, et al. 2008. J. Immunol. 180:7989. (FC IP) PubMed
- 9. Liang Y, et al. 2014. J. Immunol. 192:1277. PubMed

Description: The AF6-120.1 antibody reacts with the I-Ab MHC class II alloantigen. These class

II molecules are expressed on antigen presenting cells (including B cells) and a subset of T cells from H-2b bearing mice and are involved in antigen presentation to T cells expressing CD3/TCR and CD4 proteins. The AF6-120.1 antibody cross-reacts with H-2 κ and H-2 $^{\rm u}$ haplotypes; this antibody does not cross-react with

other haplotypes (d, f, q, r, s).

Antigen

1. Watts C. 1997. Ann. Rev. Immunol. 15:821.

References: 2. Pamer E, et al. 1998. Ann. Rev. Immunol. 16:323.