
## **Product Data Sheet**

## PerCP/Cy5.5 anti-mouse I-Ab

Catalog # / Size:	1182075 / 25 μg 1182080 / 100 μg
Clone:	AF6-120.1
Isotype:	Mouse IgG2a, κ
Immunogen:	C57BL/10J splenocytes
<b>Reactivity:</b>	Mouse
Preparation:	The antibody was purified by affinity chromatography, and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
<b>Concentration:</b>	0.2



C57BL/6 mouse splenocytes stained with AF6-120.1 PerCP/Cy5.5

## **Applications:**

Applications:	Flow Cytometry
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application. * PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.
Application Notes:	Additional reported applications (for relevant formats of this clone) include: immunohistochemical staining of frozen sections (acetone-fixed5; OCT- embedded, ethanol-fixed sections <sup>7</sup> ), immunofluorescence microscopy3 (including acetone-fixed epidermal sheets <sup>6</sup> ), immunoprecipitation <sup>7,8</sup> . Directly conjugated antibody was used for IF in (3) and (6) and for IHC in (5).
Application References:	<ol> <li>Wall KA, <i>et al.</i> 1983. <i>J. Immunol.</i> 131:1056. (FC)</li> <li>Cohn LE, <i>et al.</i> 1986. <i>P. Natl. Acad. Sci. USA</i> 83:747. (FC)</li> <li>Inaba K, <i>et al.</i> 1998. <i>J. Exp. Med.</i> 188:2163 (IF)</li> <li>Hamrah P, <i>et al.</i> 2002. <i>Invest Opthalmol Vis. Sci.</i> 43:639 (IF)</li> <li>Buono C, <i>et al.</i> 2003. <i>Arterioscler. Thromb. Vasc. Biol.</i> 23:454. (IHC)</li> <li>Wang Z, <i>et al.</i> 2004. <i>J. Immunol.</i> 172:5924. (IHC IF)</li> <li>Nakagawa TY, <i>et al.</i> 1999. <i>Immunity</i> 10:207. (IP)</li> <li>Podolin PL, <i>et al.</i> 2018. <i>J. Immunol.</i> 180:7989. (FC IP) <u>PubMed</u></li> <li>del Rio ML, <i>et al.</i> 2011. <i>Transpl. Int.</i> 24:501. (FC) <u>PubMed</u></li> <li>Weber GF, <i>et al.</i> 2014. <i>J Exp Med</i> 211:1243. <u>PubMed</u></li> </ol>
Description:	The AF6-120.1 antibody reacts with the I-Ab MHC class II alloantigen. These class II molecules are expressed on antigen presenting cells (including B cells) and a subset of T cells from H-2b bearing mice and are involved in antigen presentation to T cells expressing CD3/TCR and CD4 proteins. The AF6-120.1 antibody cross-reacts with H-2 $\kappa$ and H-2 <sup>u</sup> haplotypes; this antibody does not cross-react with other haplotypes (d, f, q, r, s).

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com 
 Antigen
 1. Watts C. 1997. Ann. Rev. Immunol. 15:821.

 References:
 2. Pamer E, et al. 1998. Ann. Rev. Immunol. 16:323.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com