
## **Product Data Sheet**

## PE anti-mouse H-2Kd

Catalog # / Size:	1183035 / 50 μg 1183040 / 200 μg	
Clone:	SF1-1.1	
Isotype:	Mouse IgG2a, к	Number
Immunogen:	BALB/c Mouse cells	
<b>Reactivity:</b>	Mouse	Relative Cell
Preparation:	The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.	Relat
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.	Ba
<b>Concentration:</b>	0.2	W



Balb/c mouse splenocytes stained with SF1-1.1 PE

## **Applications:**

Applications:	Flow Cytometry
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 0.25$ microg per $10^6$ cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.
Application Notes:	The SF1-1.1 antibody is weakly cross-reactive with H-2 $\kappa$ but does not cross-react with other haplotypes (b, j, p, q, s, v). Clone SF1-1.1 recognizes the $\alpha$ 3 domain of Kd.
	Additional reported applications (for the relevant formats) include: immunoprecipitation <sup>1,4</sup> and Western blotting2.
Application References:	<ol> <li>Noun G, <i>et al.</i> 1996. <i>J. Immunol.</i> 157:2455. (IP)</li> <li>Abasto JP, <i>et al.</i> 1993. <i>J. Immunol.</i> 151:3569. (WB)</li> <li>Bashuda H, <i>et al.</i> 1997. <i>Transplantation</i> 63:113.</li> <li>Sester M, <i>et al.</i> 2000. <i>J. Biol. Chem.</i> 34:113. (IP)</li> <li>Ma XT, <i>et al.</i> 2006. <i>Cancer Res.</i> 66:1169. (FC)</li> <li>Norian LA and Allen PM. 2004. <i>J. Immunol.</i> 173:835. (FC)</li> <li>Norian L, <i>et al.</i> 2004. <i>J. Immunol</i> 173:835. <u>PubMed</u></li> <li>Delon J, <i>et al.</i> 1998. <i>Immunity</i> 9:467.</li> </ol>
Description:	The SF1-1.1 antibody reacts with the H-2Kd MHC class I alloantigens expressed on nucleated cells from mice of the H-2Kd haplotype. H-2Kd is involved in antigen presentation to T cells expressing CD3/TCR and CD8 proteins.
Antigen References:	<ol> <li>Watts C. 1997. Annu. Rev. Immunol. 15:821.</li> <li>Pamer E, et al. 1998. Annu. Rev. Immunol. 16:323.</li> <li>York I, et al. 1996. Annu. Rev. Immunol. 14:369.</li> </ol>

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com