

Alexa Fluor® 700 anti-mouse H-2Kd

Catalog # / Size: 1183135 / 25 µg
1183140 / 100 µg

Clone: SF1-1.1

Isotype: Mouse IgG2a, κ


Immunogen: BALB/c Mouse cells

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 700 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2



BALB/c mouse splenocytes were stained with H-2Kd (clone SF1-1.1) Alexa Fluor® 700 (filled histogram) or mouse IgG2a, κ Alexa Fluor® 700 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤1.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

* Alexa Fluor® 700 has a maximum emission of 719 nm when it is excited at 633 nm / 635 nm. Prior to using Alexa Fluor® 700 conjugate for flow cytometric analysis, please verify your flow cytometer's capability of exciting and detecting the fluorochrome.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: The SF1-1.1 antibody is weakly cross-reactive with H-2k but does not cross-react with other haplotypes (b, j, p, q, s, v). Clone SF1-1.1 recognizes the α3 domain of Kd.

Additional reported applications (for the relevant formats) include: immunoprecipitation^{1,4} and Western blotting².

- Application References:**
1. Noun G, *et al.* 1996. *J. Immunol.* 157:2455. (IP)
 2. Abasto JP, *et al.* 1993. *J. Immunol.* 151:3569. (WB)
 3. Bashuda H, *et al.* 1997. *Transplantation* 63:113.
 4. Sester M, *et al.* 2000. *J. Biol. Chem.* 34:113. (IP)
 5. Ma XT, *et al.* 2006. *Cancer Res.* 66:1169. (FC)

6. Norian LA and Allen PM. 2004. *J. Immunol.* 173:835. (FC)
 7. Norian L, *et al.* 2004. *J. Immunol.* 173:835. [PubMed](#)
 8. Delon J, *et al.* 1998. *Immunity* 9:467.
-

Description: The SF1-1.1 antibody reacts with the H-2Kd MHC class I alloantigens expressed on nucleated cells from mice of the H-2Kd haplotype. H-2Kd is involved in antigen presentation to T cells expressing CD3/TCR and CD8 proteins.

Antigen
References:

1. Watts C. 1997. *Annu. Rev. Immunol.* 15:821.
2. Pamer E, *et al.* 1998. *Annu. Rev. Immunol.* 16:323.
3. York I, *et al.* 1996. *Annu. Rev. Immunol.* 14:369.