

PE/Dazzle™ 594 anti-mouse H-2Kd

Catalog # / 1183170 / 100 µg
Size: 1183165 / 25 µg

Clone: SF1-1.1

Isotype: Mouse IgG2a, κ

Immunogen: BALB/c Mouse cells


Reactivity: Mouse, Other

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide

Workshop Number: HCDM listed

Concentration: 0.2 mg/mL



BALB/c splenocytes were stained with H-2K^d (clone SF1-1.1) PE/Dazzle™ 594 (filled histogram) or mouse (SJL) IgG2a, κ PE/Dazzle™ 594 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.5 µg per million cells in 100 µL volume. It is recommended that the reagent be titrated for optimal performance for each application.

* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.

Application Notes: The SF1-1.1 antibody is weakly cross-reactive with H-2^k but does not cross-react with other haplotypes (b, j, p, q, s, v). Clone SF1-1.1 recognizes the α3 domain of K^d.

Additional reported applications (for the relevant formats) include: immunoprecipitation^{1,4} and Western blotting².

- Application References:**
1. Noun G, *et al.* 1996. *J. Immunol.* 157:2455. (IP)
 2. Abastado JP, *et al.* 1993. *J. Immunol.* 151:3569. (WB)
 3. Bashuda H, *et al.* 1997. *Transplantation* 63:113.
 4. Sester M, *et al.* 2000. *J. Biol. Chem.* 34:113. (IP)
 5. Ma XT, *et al.* 2006. *Cancer Res.* 66:1169. (FC)
 6. Norian LA and Allen PM. 2004. *J. Immunol.* 173:835. (FC)
 7. Norian L, *et al.* 2004. *J. Immunol.* 173:835. [PubMed](#)
 8. Delon J, *et al.* 1998. *Immunity* 9:467.

Description: The SF1-1.1 antibody reacts with the H-2K^d MHC class I alloantigens expressed on nucleated cells from mice of the H-2K^d haplotype. H-2K^d is involved in antigen presentation to T cells expressing CD3/TCR and CD8 proteins.

Antigen
References:

1. Watts C. 1997. *Annu. Rev. Immunol.* 15:821.
2. Pamer E, et al. 1998. *Annu. Rev. Immunol.* 16:323.
3. York I, et al. 1996. *Annu. Rev. Immunol.* 14:369.