

**APC/Cy7 anti-mouse CD11c**

**Catalog # / Size:** 1186620 / 100 µg  
1186615 / 25 µg

**Clone:** N418

**Isotype:** Hamster IgG


**Immunogen:** Mouse spleen dendritic cells

**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with APC/Cy7 under optimal conditions. The solution is free of unconjugated APC/Cy7 and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide

**Concentration:** 0.2




C57BL/6 mouse splenocytes were stained with I-A/I-E PE and CD11c (clone N418) APC/Cy7 (top) or Armenian hamster IgG APC/Cy7 isotype control (bottom).

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤1.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

**Application Notes:** Additional reported applications (for the relevant formats) include: immunoprecipitation<sup>3</sup>, immunohistochemical staining of acetone-fixed frozen sections<sup>3</sup>, and immunofluorescence microscopy<sup>5, 9</sup> (Alexa Fluor® 488 conjugated N418 was used for IHC in frozen sections<sup>10</sup>).



**Application References:**

1. Granucci F, *et al.* 1997. *J. Immunol.* 159:1794.
2. Stokes RW, *et al.* 1998. *J. Immunol.* 160:5514.
3. Metlay JP, *et al.* 1990. *J. Exp. Med.* 171:1753. (IHC, IP)
4. Ma XT, *et al.* 2006. *Cancer Research* 66:1169.
5. Chin RK, *et al.* 2006. *J. Immunol.* 177:290. (IF)
6. Cervantes-Barragan L, *et al.* 2007. *Blood* 109:1131. (FC) [PubMed](#)
7. Turnquist HR, *et al.* 2007. *J. Immunol.* 178:7018. (FC) [PubMed](#)
8. Benson MJ, *et al.* 2007. *J. Exp. Med.* doi:10.1084/jem.20070719. (FC) [PubMed](#)
9. You Y, *et al.* 2009. *J. Immunol.* 182:7343. (IF) [PubMed](#)
10. Roland CL, *et al.* 2009. *Mol. Cancer Res.* 8:1761. (IHC, FC) [PubMed](#)
11. Wikstrom M, *et al.* 2006. *J. Immunol.* 177:913. [PubMed](#)
12. Pericolini E, *et al.* 2008. *J. Leukocyte Biol.* 83:1286. [PubMed](#)
13. Randall LM, *et al.* 2008. *Infect. Immun.* 76:3312. [PubMed](#)

14. Fahlen-Yrild L, *et al.* 2009. *J. Immunol.* 183:5032. [PubMed](#)
  15. Osterholzer JJ, *et al.* 2009. *J. Immunol.* 183:8044. [PubMed](#)
  16. Bankoti J, *et al.* 2010. *Toxicol. Sci.* 115:422. (FC) [PubMed](#)
  17. Eisenach PA, *et al.* 2010. *J Cell Sci.* 123:4182. [PubMed](#)
  18. Leppin K, *et al.* 2014. *Invest. Ophthalmol. Vis. Sci.* 55:3603. [PubMed](#)
  19. Zhang K, *et al.* 2014. *PLoS Pathog.* 10:1004385. [PubMed](#)
  20. Ballet R, *et al.* 2014. *PLoS Pathog.* 10:1004550. [PubMed](#)
  21. Davis MJ, *et al.* 2015. *J Immunol.* 194:2219. [PubMed](#)
  22. Metzger DW, *et al.* 2015. *J Infect Dis.* [PubMed](#)
  23. Sharma SK, *et al.* 2015. *J Immunol.* 194:5529. [PubMed](#)
- 

**Description:** CD11c is a 150 kD glycoprotein also known as  $\alpha_X$  integrin, CR4, and p150. CD11c forms a  $\alpha_X\beta_2$  heterodimer with  $\beta_2$  integrin (CD18). It is primarily expressed on dendritic cells, NK cells, a subset of intestinal intraepithelial lymphocytes (IEL), and some activated T cells. The  $\alpha_X\beta_2$  integrin plays an important role in cell-cell contact by binding its ligands: iC3b, fibrinogen, and CD54.

**Antigen**  
**References:**

1. Barclay A, *et al.* 1997. The Leukocyte Antigen Facts Book Academic Press.
2. Springer TA. 1994. *Cell* 76:301.
3. Lopez-Rodriguez C, *et al.* 1996. *J. Immunol.* 156:3780.