


PerCP/Cyanine5.5 anti-mouse CD278 (ICOS)

Catalog # /	1187120 / 100 µg
Size:	1187115 / 25 µg
Clone:	7E.17G9
Isotype:	Rat IgG2b, κ
Immunogen:	Mouse ICOS cDNA and ICOS hexahistidine fusion protein
Reactivity:	Mouse
Preparation:	The antibody was purified by affinity chromatography and conjugated with PerCP/Cyanine5.5 under optimal conditions.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
Workshop Number:	HCDM listed
Concentration:	0.2 mg/mL



ConA-stimulated (3 days) C57BL/6 splenocytes were stained with CD3 APC and CD278 (ICOS) (clone 7E.17G9) PerCP/Cyanine5.5 (left) or rat IgG2b, κ PerCP/Cyanine5.5 isotype control (right).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 1.0 µg per million cells in 100 µL volume. It is recommended that the reagent be titrated for optimal performance for each application.

* PerCP/Cyanine5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

Application Notes: Additional reported applications (for the relevant formats) include: blocking of ligand binding.

Application References:

1. Akbari O, et al. 2002. *Nat. Med.* 8:1024.
2. Harada H, et al. 2003. *J. Clin. Invest.* 112:234.
3. McAdam AJ, et al. 2000. *J. Immunol.* 165:5035. (FC Block)
4. Tan SL, et al. 2006. *J. Immunol.* 176:2872. [PubMed](#)

Description: The 7E.17G9 antibody reacts with the 47-57 kD ICOS protein, also known as inducible costimulatory molecule, and H4. This protein is homologous to the CD28/CTLA-4 proteins. ICOS is expressed on activated T cells and a subset of thymocytes and can costimulate T cells and induce proliferation. In addition ICOS has been shown to be involved in humoral immune responses (B cell germinal center formation). The ICOS ligand, B7h/B7RP-1 and B7-H2 is constitutively expressed in B cell areas of secondary lymphoid organs and can be induced in other tissues by LPS. ICOS stimulation has been shown to potentiate TCR-mediated IL-4 and IL-10 production and has been proposed to play a role in Th2 cell development. ICOS stimulation has been shown to be involved in airway tolerance and the downregulation of pulmonary inflammation.

Antigen
References:

1. Rudd CE, et al. 2003. *Nat. Rev. Immunol.* 3:544.
2. McAdam AJ, et al. 2000. *J. Immunol.* 165:5035.
3. Mak TW, et al. 2003. *Nat. Immunol.* 4:765