

Biotin anti-mouse CD326 (Ep-CAM)

Catalog # / Size: 1191015 / 50 µg
1191020 / 500 µg

Clone: G8.8

Isotype: Rat IgG2a, κ


Immunogen: TE-71 thymic epithelial cell line

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5



Mouse thymic epithelial stromal cell line TE-71 stained with biotinylated G8.8, followed by Sav-PE

Applications:

- Applications:** Flow Cytometry
- Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.
- Application Notes:** Additional reported applications for clone G8.8 (for the relevant formats) include: immunohistochemistry of frozen sections: acetone fixed¹, with or without OCT embedding^{2,4}.
- Application References:**
1. Farr A, *et al.* 1991. *J. Histochem. Cytochem.* 39:645. (FC, IHC)
 2. Dooley J, *et al.* 2005. *J. Immunol.* 175:4331. (FC, IHC)
 3. Hinterberger M, *et al.* 2010. *Nat. Immunol.* 11:512. (FC) [PubMed](#)
 4. Gracz AD, *et al.* 2010. *Am J. Physiol Gastrointest Liver Physiol.* 298:590. (IHC) [PubMed](#)
 5. Nudel I, *et al.* 2011. *J. Immunol.* 186:891. [PubMed](#)
 6. Morimoto H, *et al.* 2012. *Biol Reprod.* 86:148. [PubMed](#)
 7. Ishii K, *et al.* 2012. *Development.* 139:1734. [PubMed](#)
 8. Takehashi M, *et al.* 2012. *Biol Reprod.* 86:178. [PubMed](#)
 9. Murakami R, *et al.* 2013. *PLoS One.* 8:73270. [PubMed](#)
 10. Taguchi K, *et al.* 2014. *Mol Cell Biol.* 34:900. [PubMed](#)
 11. Hirokawa Y, *et al.* 2014. *Am J Physiol Gastrointest Liver Physiol.* 306:547. [PubMed](#)
 12. Ding X, *et al.* 2015. *Cancer Res.* 75:330. [PubMed](#)

Description: EpCAM (CD326) mediates calcium-independent homophilic cell to cell adhesion. It may also function as a growth factor receptor. It is thought to be involved in maintaining cells in position during proliferation. Expression of EpCAM seems to correlate inversely with the level of E-cadherin (CD324). EpCAM is considered important in tumor biology.

Antigen References:

1. Borkowski TA, *et al.* 1996. *Eur. J. Immunol.* 26:110.
2. Bergsagel PL, *et al.* 1992. *J. Immunol.* 148:590.

