APC anti-mouse CD366 (Tim-3)

Catalog # / Size: 1198525 / 25 μg

1198530 / 100 µg

Clone: RMT3-23 Isotype: Rat IgG2a, κ

Reactivity: Mouse

Preparation: The antibody was purified by affinity


chromatography and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and

unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

Concentration: 0.2



Mouse Tim-3 transfected cells were stained with anti-mouse CD366 (Tim-3, clone RMT3-23) APC (filled histogram) or rat IgG2a, κ APC isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended E

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is <0.25 microg per million cells in 100 microl, volume. It is

this reagent is \leq 0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each

application.

Application

Notes:

Additional reported applications (for relevant formats) include: *in vitro*1 and *in vivo*2 blocking of Tim-3, and immunohistochemical staining of frozen sections2. The LEAF $^{\text{TM}}$ purified antibody (Endotoxin EU/microg, Azide-Free, 0.2 μ m filtered) is

recommended for functional assays (Cat. No. 119707).

Application References:

1. Nakae S, et al. 2007. Blood 110(7):2565-8. (FC, Block)

2. Oikawa T, et al. 2006. J. Immunol. 177(7):4281-7. (FC, Block, IHC)

Description:

CD366 (Tim-3) is a transmembrane protein also known as T cell immunoglobulin and mucin domain containing protein-3. Tim-3 is expressed at high levels on Th1 lymphocytes and CD11b⁺ macrophages. Tim-3 has also been shown to exist as a soluble protein. Cells expressing Tim-3 are present at high levels in the CNS of animals at the onset of experimental autoimmune encephalomyelitis (EAE), a disease mediated by lymphocytes secreting Th1-like cytokines. Tim-3 has been proposed to inhibit Th1-mediated immune responses and promote immunological tolerance.

Antigen

1. Sabatos CA, et al. 2003. Nat. Immunol. 4:1102.

References: 2. Sanchez-Fueyo A, et al. 2003. Nat. Immunol. 4:1102.

3. Kuchroo VK, et al. 2003. Nat. Rev. Immunol. 3:454.

4. Mooney L, et al. <